(19) World Intellectual Property Organization International Bureau





PCT

(43) International Publication Date 16 November 2006 (16.11.2006)

(51) International Patent Classification:

A61K 31/551 (2006.01)

(21) International Application Number: PCT/US2006/017677

(22) International Filing Date: 5 May 2006 (05.05.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: 60/678,667 5 May 2005 (05.05.2005)

(71) Applicant (for all designated States except US): VALEANT RESEARCH & DEVELOPMENT [US/US]; 3300 Hyland Avenue, Costa Mesa, CA 92626 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): GIRARDET, Jean, Luc [FR/US]; 3300 Hyland Avenue, Costa Mesa, CA 92626 (US). KOH, Yung-hyo [KR/US]; 22 Oakhurst Road, Irvine, CA 92620 (US). SHAW, Stephanie [US/US]; 2684 Pepperdale Drive, Rowland Heights, CA 91748 (US). KIN, Hong Woo [KR/US]; 400 Santa Maria Drive, Irvine, CA 92608 (US).

(10) International Publication Number $WO\ 2006/122003\ A2$

- (74) Agent: REESE, Heidi, L.; BROWN RAYSMAN MILL-STEIN FELDER & STEINER LLP, 900 Third Avenue, New York, NY 10022 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, IP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NΛ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

 without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DIARYL-PURINE, AZAPURINES AND -DEAZAPURINES AS NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS FOR TREATMENT OF HIV

(57) Abstract: This application concerns certain 2-phenylamino-6-aryl amino-, 6-aryloxy-, and 6- arylthio- purines, -azapurines and -deazapurines. These compounds are non-nucleoside reverse transcriptase inhibitors and have potential as anti-HIV treatment.

WO 2006/122003 PCT/US2006/017677

DIARYL-PURINES, -AZAPURINES AND -DEAZAPURINES AS NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS FOR TREATMENT OF HIV

Cross-reference to Related Applications

This application claims priority to U.S. Provisional Application Ser. No. 60/678,667, filed May 5, 2005, the entirety of which is incorporated herein by reference.

Field of the Invention

This application concerns certain 2-phenylamino-6-aryl amino-, 6-aryloxy-, and 6-arylthio- purines, -azapurines and -deazapurines. These compounds are non-nucleoside reverse transcriptase inhibitors and have potential as anti-HIV treatment.

Background of the Invention

Human Immunodeficiency Virus (HIV) presents a public-health and social catastrophe too well known to require documentation. One therapeutic approach to HIV has been inhibition of the viral RNA-dependent RNA polymerase; this enzyme is frequently referred to as "reverse transcriptase," abbreviated "RT." The first RT inhibitors were nucleoside analogs such as AZT and ddI. Although such nucleoside RT inhibitors were frequently effective against the wild-type virus, any single-drug treatment has been hobbled by the virus's ability to readily produce drug-resistant mutants. This has led to an intense search for non-nucleoside RT inhibitors ("NNRTIs") which are both effective and capable of retaining their effectiveness despite drug-resistance mutations. A recent review of NNRTIs can be found Balzarni, J., 2004, Cur. Top. Med. Chem. 4, 921-44 (Erratum ibid. 4, 1825).

Four leading NNRTI are: 1) Efavirenz (4S)-6-chloro-4-(cyclopropylethynyl)-1,4-dihydro-4-(trifluoromethyl)-2H-3,1-benzoxazin-2-one; 2) Capravirine: 1H-Imidazole-2-methanol, 5-((3,5-dichlorophenyl)thio)-4-(1-methylethyl)-1-(4-pyridinylmethyl)-carbamate (ester); 3) Etravirine (TMC 125): 4-((6-amino-5-bromo-2-((4-cyanophenyl)amino)-4-pyrimidinyl)oxy)-3,5-dimethyl-benzonitrile; and 4) Rilpivirine (TMC-278): 4-([4-[(4-[(1E)-2-cyanoethenyl]-2,6-dimethyl-phenyl)amino]-2-pyrimidinyl)amino]benzonitrile. Rilpivirine and Etravirine belong to a subclass of NNRTIs called diarylpyrmidines ("DAPY"). For a review of these DAPY NNRTIs see Ludovici, D.W., et al., 2002, Bioorg. Med. Chem. Lett.

11, 2235-9. An extensive patent literature also exists for DAPY. U.S. Patent No. 6,197,779; WO 00/27850; WO 2003/016306; and WO 2004/069812, all assigned to Janssen Pharmaceuticals.

Diaryl compounds similar to Etravirine and Rilpivirine where the pyrimidine moiety is replaced by a purine are described in WO 2005/028479, which also is assigned to Janssen.

Brief Description of the Invention

The invention provides a compound of formula I

I

where the dashed line represents a double bond that may be located either between A and B or between B and D,

where A is -N=, N(Z) or C(Z);

B is CH or =N-;

D is CW or =N-, or N(W);

15 T is NH, O or S;

Z is H, F, Cl, Br, CH₃, CH₂CH₃, cyclopropyl, or benzyl, in which the phenyl moiety of the benzyl group is optionally substituted with methyl or methoxy, provided that Z is not F or Cl when A is NZ;

W is H, F, Cl, Br, methyl, ethyl, cyclopropyl, allyl, CH₂CF₃, cyanomethyl, cyanoethyl,

20 CH=CHCN, or benzyl, in which the phenyl moiety of the benzyl group is optionally substituted with one or two groups selected independently from methoxy and methyl, provided that W is not F or Cl when D is NW;

V is F, Cl, CN, SO₂CH₃, SO₂NH₂, SO₂NHCH₃, C≡CCH₃, or CH=CHCN; provided that when D is CW, A is not CZ and further provided that when neither A nor D is

25 CZ or CW, then B is CH; and

Ar is selected from (a), (b), (c), and (d) below:

(a)
$$R^6$$
 R^5

(b)
$$R^6$$
 Q Q' R^4 R^5

(d)
$$R^{6}$$
 R^{8} R^{10} R^{11}

wherein each R^P is selected from among methyl, ethyl, propyl, isopropyl, cyclopropylmethyl, or C_3 - C_6 cycloalkyl, cyano, CH=CHCN, Cl, Br, I, acetyl and alkylamino; R^4 , R^5 , and each R^6 are independently selected from among H, F, Cl, Br, CH₃, CF₃, CH₂F, CHF₂, isopropyl, cyclopropyl, OCH₃, OH, OCF₃, NH₂ and NHCH₃, or R^6 and R^P on adjacent ring atoms, together with the ring atoms to which they are attached, form an additional fused five-membered ring; Q and Q' are independently selected from N and CH; R^7 is Cl, Br, I, CH₃, CF₃, OCH₃, isopropyl, cyclopropyl, *t*-butyl, or cyclobutyl; and $R^8 - R^{11}$ are, independently, H or CH₃.

Compounds of formula I have inhibitory activity against both wild-type and mutated forms of human immunodeficiency virus type 1 (HIV-1).

15

10

Detailed Description of the Invention

In one embodiment this invention provides a compound of formula IA, in which the 6-linker T of formula I is T', which may be O or S.

5

10

25

When T of formula I or T' of formula IA is O, the invention excludes 1) compounds where both R^p and V are CH=CHCN or cyano unless at least one of A or D is neither -N= nor - NH- and 2) compounds where a) R is CH=CHCN, cyano, or methyl; b) V is cyano or CH=CHCN; and c) A and D are both one of -N=; N-benzyl or N-(substituted benzyl).

In one subgeneric embodiment, the invention provides a compound of formula IA
where Ar is selected from 4-cyclopropyl phenyl; 4-cyclopropylmethyl phenyl; 4bromophenyl; 4-cyclopropyl-naphth-1-yl; 2,6-dimethyl-4-cyanophenyl; 2,6-dimethoxy-4cyanophenyl; 2,6-dimethyl-4-(2-cyanoethenyl) phenyl; 2,6-dimethoxy-4-(2-cyanoethenyl)
phenyl; 2-methyl-4-cyclopropyl phenyl; 2,6-dimethyl-4-cyclopropyl phenyl; 2,6-ditrifluoromethyl-4-cyclopropyl phenyl; 2,4,6-trimethyl phenyl; and 2,6-dimethyl-4-acetyl
phenyl.

In another subgeneric embodiment, the invention contemplates a compound of formula IA where Ar is selected from the following: 5-cyclopropyl-8-quinolyl; 5-isopropyl-8-quinolyl; 5-cyclopropyl-7-trifluoromethyl-8-quinolyl; 5-acetyl-8-quinolyl; 5-cyano-7-methoxy-8-quinolyl; 5-cyano-7-methyl-8-quinolyl; 5-cyclopropyl-7-trifluoromethoxy-8-isoquinolyl; 5-cyano-8-isoquinolyl; 5-cyano-7-methoxy-8-isoquinolyl; 5-cyano-7-methyl-8-isoquinolyl; 5-cyclopropyl-7-methyl-8-cinnolyl; 3-cyclopropyl-7-methyl-8-cinnolyl; 3-cyclopropyl-7-methyl-8-cinnolyl;

In another subgeneric embodiment, the invention provides a compound of formula IA-1

5

10

15

20

where Ar, V, W, and Z are defined as for formula I.

In another subgeneric embodiment, the invention provides a compound of formula IA-2

IA-2 where Ar, V, W, and Z are defined as for formula I.

In another subgeneric embodiment, the invention provides a compound of formula IA-3

where Ar, V, W, and Z are defined as for formula I.

In another subgeneric embodiment, this invention provides a compound of formula IA-4

where Ar, V, W, and Z are defined as for formula I.

In another subgeneric embodiment, this invention provides a compound of formula 5 IA-5

IA-5

10

where Ar, V, W, and Z are defined as for formula I.

In another subgeneric embodiment, this invention provides a compound of formula IA-6

15

IA-6

where Ar, V, W, and Z are defined as for formula I.

20

In another subgeneric embodiment, this invention provides a compound of formula IA-7

IA-7

25 where Ar, V, W, and Z are defined as for formula I.

In another subgeneric embodiment, this invention provides a compound of formula IA-8

5

10

20

where Ar, V, W, and Z are defined as for formula I.

In another subgeneric embodiment, this invention provides a compound of formula IA-9

IA-9

where Ar, V, W, and Z are defined as for formula I.

In another subgeneric embodiment, this invention provides a compound of formula 15 IA-10

IA-10

where Ar, V, W, and Z are defined as for formula I.

In another embodiment, this invention provides a compound of formula IB

IB

where all substituents are as described above, except that when Ar is (c), this invention excludes compounds in which V is either cyano or CH=CHCN, unless A or D is CZ or CW.

In one subgeneric embodiment, the invention provides a compound of formula IB where Ar is (c), subject to the exclusion in the immediately preceding paragraph.

In a more specific subgeneric embodiment, the invention provides a compound of formula IB where Ar is

$$\mathbb{R}^{\theta}$$
 \mathbb{R}^{7}

5

where R^p is CN, CH=CHCN, or cyclopropyl; where R⁶ and R⁷ are either both methyl or both methoxy; and subject to the exclusion described above for formula IB.

In another subgeneric embodiment, this invention provides a compound of formula IB-1.

10

IB-1

In another subgeneric embodiment, this invention provides a compound of formula IB-2.

15

where Ar, V, W, and Z are as described above for formula IB.

In another subgeneric embodiment, the invention provides a compound of formula IB-3.

IB-3

where Ar, W, and Z are as described above for formula IB.

In another subgeneric embodiment, the invention provides a compound of formula IB-4.

5

IB-4

where Ar, V, and Z are as described above for formula IB.

In more specific embodiments, the invention provides compounds of any of IA-1, IA-2, IA-3, IA-4, IA-5, IA-6, IA-7, IA-8, IA-9, IA-10, IB-1, IB-2, IB-3, and IB-4, where

10 Ar is (a).

In additional more specific embodiments, the invention provides compounds of any of IA-1, IA-2, IA-3, IA-4, IA-5, IA-6, IA-7, IA-8, IA-9, IA-10, IB-1, IB-2, IB-3, and IB-4, where Ar is (b).

In additional more specific embodiments, the invention provides compounds of any of IA-1, IA-2, IA-3, IA-4, IA-5, IA-6, IA-7, IA-8, IA-9, IA-10, IB-1, IB-2, IB-3, and IB-4, where Ar is (c).

In additional more specific embodiments, the invention provides compounds of any of IA-1, IA-2, IA-3, IA-4, IA-5, IA-6, IA-7, IA-8, IA-9, IA-10, IB-1, IB-2, IB-3, and IB-4, where Ar is (d).

In a more specific subgeneric embodiment, this invention provides or contemplates a compound of formula IA-7, IA-8, IA-9, or IA-10, where Ar is 4-cyclopropyl-, 4-acetyl-, 4-methyl-, 4-bromo-, or 4-cyano-2,6-di-substituted phenyl.

In another more specific subgeneric embodiment, this invention provides or contemplates a compound of formula IA-1, IA-2, IA-3, or IA-4, where Ar is 4-cyclopropyl-, 4-acetyl-, 4-methyl-, 4-bromo-, or 4-cyano-2,6-di-substituted phenyl.

In another more specific subgeneric embodiment, this invention provides or contemplates a compound of formula IA-5 or IA-6, where Ar is 4-cyclopropyl-, 4-acetyl-, 4-methyl-, 4-bromo-or 4-cyano-2,6-di-substituted phenyl.

Synthetic procedures

Compounds of this invention which are of the **7-deaza-8-azapurine** type can be prepared according to Scheme 1.

Scheme 1

$$NH_2$$
 NNH
 CH_3I
 NNH
 CN
 NNH
 CN
 NNH
 NNH

10

15

5

Compound (1), 2-mercapto-6-hydroxy-7-deaza-8-aza-purine, can be synthesized by published procedures known to those skilled in the art. Youssif, S., et al., 2003, Bull. Kor. Chem. Soc., 24, 1429-32; Bontems, R.J., et al., 1990, J. Med. Chem. 33, 2174-8; Badger, G.M., & Rao, R.P., 1965, Aust. J. Chem. 18, 1267-71.

Alternatively, the 7-deaza-8-azapurines can be synthesized according to Scheme 2, where "PMBCI" is *p*-methoxy benzyl chloride. The starting material is prepared by published procedures known to those skilled in the art. Seela, F., 1999, *Helv. Chim. Act.* 82, 105-124; Taylor, E., 1992, *Tetrahedron* 48, 8089-100; Seela, F., 1986, *Helv. Chim. Act.* 69, 1602-1613.

Scheme 2

$$N-NH$$
 CI
 $N-NH$
 CI
 $N-NH$
 CI
 $N-NH$
 $N-N$

The 8-aza-9-deazapurines of this invention can be synthesized according to Scheme 3. The synthesis of the starting material was described by Lewis, A.F., & Townsend, L.B., 1982, *J. Am. Chem. Soc.* 104, 1073-78.

The 9-deazapurines of this invention can be synthesized by Scheme 4. The synthesis of the starting material is described by Kielich, Klaus, *ed.*, "Synthetic Communications" **2002** vol. 32, pp-3797-3802.

Scheme 4

5

The 7-deazapurines of this invention are prepared by the procedure of Scheme 5. The starting material can be synthesized by the condensation of 2,6-diamino-1,2-dihydro[3H]pyrimidin-4-one with chloroacetaldehyde followed by treatment with phosphorus oxychloride, as indicated in Examples 1 and 3.

Bn
$$\frac{NH_2}{TFA, CF_3CH_2OH, 100^{\circ}C}$$
 $\frac{NH_2}{TFA, CF_3CH_2OH, 100^{\circ}C}$ $\frac{NH_2$

The **purine** compounds of this invention can be synthesized by strategies similar to those provided above, using N⁷-benzyl-2,6-dichloropurine as the starting material. This procedure is illustrated in WO 2005/028479.

5 Example 1

Step Al:

10

15

CI + HN NH₂ NaOAc NH NH₂ Nh
$$_2$$
 NaOAc NH NH₂ Nh $_2$ NaOAc NH NH $_2$

2-Amino-3,7-dihydro-pyrrolo[2,3-d]pyrimidin-4-one. To a mixture of 2,4-diamino-6-hydroxypyrimidine (20.0 g, 159 mmol) and NaOAc (26.0 g, 317 mmol) in H₂O (300 mL) at 65°C was added a solution of chloroacetaldehyde (22.0 mL, 50% in H₂O, 173 mmol) in H₂O (22 mL) dropwise for 90 min. The mixture was stirred at 65°C for an additional 2 h and cooled to room temperature. The reaction mixture was concentrated *in vacuo* to one third of its original volume and stored at 4 °C for 16 h. The light pink precipitates were filtered, washed with an ice cold H₂O (5 mL), and dried under high vacuum for 16 h. The precipitates were placed in Soxhlet extractor and refluxed with methanol (200 mL) for 24 h. The methanol was concentrated to give 13.3 g (56%) of 2-amino-3,7-dihydro-pyrrolo[2,3-d]pyrimidin-4-one as a light pink solid.

Step A2:

5

10

15

20

25

4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine. To a solution of 2-amino-3,7-dihydro-pyrrolo[2,3-d]pyrimidin-4-one (5.00 g, 33.3 mmol), dimethylaniline (4.22 mL, 41.0 mmol) and benzyltriethylammonium chloride (15.2 g, 66.6 mmol) in acetonitrile (25 mL) at room temperature under argon was added POCl₃ (18.6 mL, 200 mmol) dropwise for 30 min. The mixture was refluxed at 85°C for 3 h and cooled to room temperature. The reaction was concentrated *in vacuo* to brown oil and to the oil was added an ice cold H₂O (10 mL). The pH of the solution was adjusted to 5 by the addition of an aqueous NH₄OH solution. Silica gel chromatography (CH₂Cl₂:MeOH = 95: 5) yielded 2.53 g (45%) of 4-chloro-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine as a light yellow solid. The product was then benzylated at N⁷ using standard techniques.

Step C:

7-benzyl-4-(2,4,6-trimethyl-phenoxy)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine. To a solution of 2,4,6-trimethylphenol (161 mg, 1.16 mmol) in 1-methyl-2-pyrridone (2 mL) in a sealed tube was added NaH (46 mg, 1.16 mmol). The reaction mixture was stirred at room temperature for 15 min and a solution of 7-benzyl-4-chloro-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (100 mg, 0.39 mmol) in 1-methyl-2-pyrridone (1 mL) was added to the mixture. The mixture was heated at 150 °C for 16 h and cooled to room temperature. The reaction mixture was poured into ice water and extracted with EtOAc (2 x 20 mL). The combined organic solution was washed with H₂O (20 mL) and brine (20 mL), dried with Na₂SO₄, and

concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 75:25) yielded 107 mg (77%) of 7-benzyl-4-(2,4,6-trimethyl-phenoxy)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine.

Step D:

5

10

15

7-benzyl-2-fluoro-4-(2,4,6-trimethyl-phenoxy)-7H-pyrrolo[2,3-d]pyrimidine. To 7-benzyl-4-(2,4,6-trimethyl-phenoxy)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (105 mg, 0.29 mmol) in a polyethylene flask at -50 °C under argon was added 60% HF in pyridine (12 mL). To the resulting solution tert-butylnitrite (0.052 mL, 0.44 mmol) was added dropwise for 5 min. The reaction was warmed to -40 °C and stirred for 30 min at the temperature. The reaction mixture was diluted with CHCl₃ (100 mL) and poured into K₂CO₃ (3 g) in a beaker. Ice water (50 mL) was carefully added to the mixture. The CHCl₃ layer was separated, washed with aqueous NaHCO₃ solution (20 mL) and brine (20 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 75:25) yielded 72 mg (68%) of 7-benzyl-2-fluoro-4-(2,4,6-trimethyl-phenoxy)-7H-pyrrolo[2,3-d]pyrimidine as a light yellow solid.

Step E:

4-[7-benzyl-4-(2,4,6-trimethyl-phenoxy)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamino]benzonitrile. To a solution of 4-aminobenzonitrile (101 mg, 0.86 mmol) in 1-methyl-2pyrridone (1 mL) was added NaH (34 mg, 0.86 mmol). The reaction mixture was stirred at room temperature for 15 min and a solution of 7-benzyl-2-fluoro-4-(2,4,6-trimethyl-phenoxy)-7H-pyrrolo[2,3-d]pyrimidine (62 mg, 0.17 mmol) in 1-methyl-2-pyrridone (1 mL) was added to the mixture. The mixture was stirred at room temperature for 1 h, poured into ice water, and extracted with EtOAc (2 x 20 mL). The combined organic solution was washed with H_2O (20 mL) and brine (20 mL), dried with Na_2SO_4 , and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 75:25) yielded 64 mg (82%) of 4-[7-benzyl-4-(2,4,6-trimethyl-phenoxy)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamino]-benzonitrile.

Step F:

4-[4-(2,4,6-Trimethyl-phenoxy)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamino]-benzonitrile.

To a solution of 4-[7-benzyl-4-(2,4,6-trimethyl-phenoxy)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamino]-benzonitrile (38 mg, 0.083 mmol) in 1,2-dichlorobenzene (1 mL) was added aluminum chloride (55 mg, 0.42 mmol). The reaction mixture was stirred at 160 °C for 4 h and cooled to room temperature. The mixture was poured into ice water and extracted with CH_2Cl_2 (2 x 10 mL). The combined organic solution was washed with brine (10 mL), dried with Na_2SO_4 , and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 50:50) yielded 15 mg (49%) of 4-[4-(2,4,6-trimethyl-phenoxy)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamino]-benzonitrile as a tan solid.

20

Example 2

5 Step A:

10

7-benzyl-4-(2,4,6-trimethyl-phenylsulfanyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine. To a solution of 2,4,6-trimethylbenzene-1-thiol (231 mg, 1.52 mmol) in 1-methyl-2-pyrridone (2 mL) was added NaH (58 mg, 1.52 mmol). The reaction mixture was stirred at room temperature for 15 min and a solution of 7-benzyl-4-chloro-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (131 mg, 0.51 mmol) in 1-methyl-2-pyrridone (2 mL) was added to the mixture. The mixture was heated at 60 °C for 16 h and cooled to room temperature. The reaction was poured into ice water and extracted with EtOAc (2 x 20 mL). The combined organic solution

was washed with H₂O (20 mL) and brine (20 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 75:25) yielded 180 mg (94%) of 7-benzyl-4-(2,4,6-trimethyl-phenylsulfanyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine.

Step B:

5

10

15

7-benzyl-2-fluoro-4-(2,4,6-trimethyl-phenylsulfanyl)-7H-pyrrolo[2,3-d]pyrimidine. To 7-benzyl-4-(2,4,6-trimethyl-phenylsulfanyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (155 mg, 0.41 mmol) in a polyethylene flask at -50 °C under argon was added 60% HF in pyridine (12 mL). To the solution was added tert-butylnitrite (0.074 mL, 0.62 mmol) dropwise for 5 min. The reaction was warmed to -40 °C and stirred for 30 min at the temperature. The reaction was diluted with CHCl₃ (100 mL) and poured into K₂CO₃ (3 g) in a beaker. To the mixture was carefully added ice water (50 mL). The CHCl₃ layer was separated, washed with aqueous NaHCO₃ solution (20 mL) and brine (20 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 75:25) yielded 118 mg (77%) of 7-benzyl-2-fluoro-4-(2,4,6-trimethyl-phenylsulfanyl)-7H-pyrrolo[2,3-d]pyrimidine as a yellow solid.

Step C:

4-[7-benzyl-4-(2,4,6-trimethyl-phenylsulfanyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamino]-

benzonitrile. To a solution of 4-aminobenzonitrile (184 mg, 1.56 mmol) in 1-methyl-2-pyrridone (2 mL) was added NaH (62 mg, 1.56 mmol). The reaction mixture was stirred at room temperature for 15 min and a solution of 7-benzyl-2-fluoro-4-(2,4,6-trimethyl-phenylsulfanyl)-7H-pyrrolo[2,3-d]pyrimidine (118 mg, 0.31 mmol) in 1-methyl-2-pyrridone (2 mL) was added to the mixture. The mixture was stirred at room temperature for 4 h, then

poured into ice water and extracted with EtOAc (2 x 20 mL). The combined organic solution was washed with H₂O (20 mL) and brine (20 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 75:25) yielded 123 mg (83%) of 4-[7-benzyl-4-(2,4,6-trimethyl-phenylsulfanyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamino]-benzonitrile.

Step D:

5

15

$\hbox{$4-[4-(2,4,6-Trimethyl-phenylsulfanyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamino]-$}$

benzonitrile. To a solution of 4-[7-benzyl-4-(2,4,6-trimethyl-phenylsulfanyl)-7H
pyrrolo[2,3-d]pyrimidin-2-ylamino]-benzonitrile (103 mg, 0.21 mmol) in 1,2dichlorobenzene (2 mL) was added aluminum chloride (87 mg, 0.65 mmol). The reaction mixture was stirred at 160 °C for 1.5 h and cooled to room temperature. The mixture was poured into ice water and extracted with CH₂Cl₂ (2 x 10 mL). The combined organic solution was washed with brine (10 mL), dried with Na₂SO₄, and concentrated to dryness.

Silica gel chromatography (Hexanes:EtOAc = 50:50) yielded 28 mg (34%) of 4-[4-(2,4,6-trimethyl-phenylsulfanyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamino]-benzonitrile as a tan solid.

Example 3

5 Steps A and B as in Example 1. Step C:

15

10 4-(2-Amino-7-benzyl-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy)-3,5-dimethyl-benzonitrile.

To a solution of 4-hydroxy-3,5-dimethylbenzonitrile (1.62 mg, 11.0 mmol) in 1-methyl-2-pyrridone (5 mL) in a sealed tube was added NaH (441 mg, 11.0 mmol). The reaction mixture was stirred at room temperature for 15 min and a solution of 7-benzyl-4-chloro-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (950 mg, 3.67 mmol) in 1-methyl-2-pyrridone (5 mL) was added to the mixture. The mixture was heated at 150 °C for 16 h and cooled to room temperature. The reaction was poured into ice water and extracted with EtOAc (2 x 50 mL). The combined organic solution was washed with H_2O (50 mL) and brine (50 mL), dried with Na_2SO_4 , and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 75:25)

5 yielded 1.12 mg (83%) of 4-(2-amino-7-benzyl-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy)-3,5-dimethyl-benzonitrile.

Step D:

4-(7-benzyl-2-fluoro-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy)-3,5-dimethyl-benzonitrile. To
4-(2-amino-7-benzyl-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy)-3,5-dimethyl-benzonitrile (70 mg,
0.19 mmol) in a polyethylene flask at -50 °C under argon was added 60% HF in pyridine (12 mL). To the solution was added tert-butylnitrite (0.068 mL, 0.57 mmol) dropwise for 5 min.

The reaction was warmed to -40 °C and stirred for 30 min at the temperature. The reaction was diluted with CHCl₃ (100 mL) and poured into K₂CO₃ (3 g) in a beaker. To the mixture

was carefully added ice water (50 mL). The CHCl₃ layer was separated, washed with aqueous NaHCO₃ solution (20 mL) and brine (20 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 75:25) yielded 36 mg (51%) of 4(7-benzyl-2-fluoro-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy)-3,5-dimethyl-benzonitrile.

Step E:

4-[7-benzyl-2-(4-cyano-phenylamino)-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy]-3,5-dimethyl-benzonitrile. To a solution of 4-aminobenzonitrile (54 mg, 0.46 mmol) in 1-methyl-2-pyrridone (1 mL) was added NaH (18 mg, 0.46 mmol). The reaction mixture was stirred at room temperature for 15 min and a solution of 4-(7-benzyl-2-fluoro-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy)-3,5-dimethyl-benzonitrile (34 mg, 0.091 mmol) in 1-methyl-2-pyrridone (1 mL) was added to the mixture. The mixture was stirred at room temperature for 1 h, poured into ice water, and extracted with EtOAc (2 x 20 mL). The combined organic solution was washed with H₂O (20 mL) and brine (20 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 75:25) yielded 28 mg (65%) of 4-[7-benzyl-2-(4-cyano-phenylamino)-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy]-3,5-dimethyl-benzonitrile.

Step F:

15

20

10

4-[2-(4-Cyano-phenylamino)-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy]-3,5-dimethyl-benzonitrile. To a solution of 4-[7-benzyl-2-(4-cyano-phenylamino)-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy]-3,5-dimethyl-benzonitrile (28 mg, 0.060 mmol) in 1,2-dichlorobenzene (1 mL) was added aluminum chloride (40 mg, 0.30 mmol). The reaction mixture was stirred at 160 °C for 45 min and cooled to room temperature. The mixture was poured into ice water and extracted with CH_2Cl_2 (2 x 10 mL). The combined organic solution was washed with brine (10 mL), dried with Na_2SO_4 , and concentrated to dryness. Silica gel chromatography

(Hexanes:EtOAc = 50:50) yielded 6 mg (27%) of 4-[2-(4-cyano-phenylamino)-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy]-3,5-dimethyl-benzonitrile as a tan solid.

Example 4

5

Step A:

10

15

7-benzyl-N4-(2,4,6-trimethyl-phenyl)-7H-pyrrolo[2,3-d]pyrimidine-2,4-diamine. To a suspension of 7-benzyl-4-chloro-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (200 mg, 0.78 mmol) and 2,4,6-trimethylaniline (0.44 mL, 3.08 mmol) in 2,2,2-trifluoroethanol (4 mL) was added trifluoroacetic acid (0.48 mL, 6.24 mmol). The resulting solution was heated at 100 °C for 2 days and cooled to room temperature. The reaction was concentrated to brown oil and diluted with CH_2Cl_2 (30 mL). The organic solution was washed with aqueous NaHCO₃

solution (20 mL) and brine (20 mL), dried with Na_2SO_4 , and concentrated to dryness. Silica gel chromatography (CH₂Cl₂:MeOH = 95:5) yielded 251 mg (90%) of 7-benzyl-N4-(2,4,6-trimethyl-phenyl)-7H-pyrrolo[2,3-d]pyrimidine-2,4-diamine.

Step B:

5

10

15

(7-benzyl-2-fluoro-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-(2,4,6-trimethyl-phenyl)-amine. To 7-benzyl-N4-(2,4,6-trimethyl-phenyl)-7H-pyrrolo[2,3-d]pyrimidine-2,4-diamine (251 mg, 0.70 mmol) in a polyethylene flask at -50 °C under argon was added 60% HF in pyridine (24 mL). To the solution was added tert-butylnitrite (0.42 mL, 3.5 mmol) dropwise for 10 min. The reaction was warmed to -40 °C and stirred for 30 min at the temperature. The reaction was diluted with CHCl₃ (200 mL) and poured into K₂CO₃ (6 g) in a beaker. To the mixture was carefully added ice water (100 mL). The CHCl₃ layer was separated, washed with aqueous NaHCO₃ solution (40 mL) and brine (40 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 75:25) yielded 56 mg (22%) of (7-benzyl-2-fluoro-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-(2,4,6-trimethyl-phenyl)-amine.

Step C:

4-[7-benzyl-4-(2,4,6-trimethyl-phenylamino)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamino]-benzonitrile. To a suspension of (7-benzyl-2-fluoro-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-(2,4,6-trimethyl-phenyl)-amine (42 mg, 0.12 mmol) and 4-aminobenzonitrile (55 mg, 0.47 mmol) in 2,2,2-trifluoroethanol (4 mL) was added trifluoroacetic acid (0.072 mL, 0.94 mmol). The resulting solution was heated at 90 °C for 16 h, then cooled to room temperature.

The reaction was concentrated to produce a brown oil and diluted with CH_2Cl_2 (30 mL). The organic solution was washed with H_2O (20 mL) and brine (20 mL), dried with Na_2SO_4 , and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 75:25) yielded 34 mg (64%) of 4-[7-benzyl-4-(2,4,6-trimethyl-phenylamino)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamino]-benzonitrile.

Step D:

10

5

4-[4-(2,4,6-Trimethyl-phenylamino)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamino]-

benzonitrile. To a solution of 4-[7-benzyl-4-(2,4,6-trimethyl-phenylamino)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamino]-benzonitrile (34 mg, 0.074 mmol) in 1,2-dichlorobenzene (1 mL) was added aluminum chloride (50 mg, 0.37 mmol). The reaction mixture was stirred at 160 °C for 2 h and cooled to room temperature. The mixture was poured into ice water and extracted with CHCl₃ (2 x 10 mL). The combined organic solution was washed with brine (10 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (CH₂Cl₂:Acetone = 90:10) yielded 5 mg (19%) of 4-[4-(2,4,6-trimethyl-phenylamino)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamino]-benzonitrile as a tan solid.

20

15

Example 5

5 Step A:

10

4-(2-Amino-7-benzyl-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy)-3,5-dimethyl-

benzonitrile. To a solution of 4-hydroxy-3,5-dimethylbenzonitrile (1.62 mg, 11.0 mmol) in 1-methyl-2-pyrridone (5 mL) in a sealed tube was added NaH (441 mg, 11.0 mmol). The reaction mixture was stirred at room temperature for 15 min and a solution of 7-benzyl-4-chloro-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (950 mg, 3.67 mmol) in 1-methyl-2-pyrridone (5 mL) was added to the mixture. The mixture was heated at 150 °C for 16 h and cooled to

room temperature. The reaction was poured into ice water and extracted with EtOAc (2 x 50 mL). The combined organic solution was washed with H₂O (50 mL) and brine (50 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 75:25) yielded 1.12 mg (83%) of 4-(2-amino-7-benzyl-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy)-3,5-dimethyl-benzonitrile.

Step B:

10

5

4-(7-benzyl-2-fluoro-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy)-3,5-dimethyl-benzonitrile. To 4-(2-amino-7-benzyl-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy)-3,5-dimethyl-benzonitrile (70 mg, 0.19 mmol) in a polyethylene flask at -50 °C under argon was added 60% HF in pyridine (12 mL). To the solution was added tert-butylnitrite (0.068 mL, 0.57 mmol) dropwise for 5 min.
15 The reaction was warmed to -40 °C and stirred for 30 min at the temperature. The reaction was then diluted with CHCl₃ (100 mL) and poured into K₂CO₃ (3 g) in a beaker. Ice water (50 mL) was carefully added. The CHCl₃ layer was separated, washed with aqueous NaHCO₃ solution (20 mL) and brine (20 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 75:25) yielded 36 mg (51%) of 4-(7-benzyl-2-fluoro-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy)-3,5-dimethyl-benzonitrile.

Step C:

4-[7-benzyl-2-(4-cyano-phenylamino)-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy]-3,5-dimethyl-benzonitrile. To a solution of 4-aminobenzonitrile (54 mg, 0.46 mmol) in 1-methyl-2-pyrridone (1 mL) was added NaH (18 mg, 0.46 mmol). The reaction mixture was stirred at room temperature for 15 min and a solution of 4-(7-benzyl-2-fluoro-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy)-3,5-dimethyl-benzonitrile (34 mg, 0.091 mmol) in 1-methyl-2-pyrridone (1 mL) was added to the mixture. The mixture was stirred at room temperature for 1 h, poured into ice water and extracted with EtOAc (2 x 20 mL). The combined organic solution was washed with H₂O (20 mL) and brine (20 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 75:25) yielded 28 mg (65%) of 4-[7-benzyl-2-(4-cyano-phenylamino)-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy]-3,5-dimethyl-benzonitrile.

Step D:

10

4-[2-(4-Cyano-phenylamino)-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy]-3,5-dimethyl-benzonitrile. To a solution of 4-[7-benzyl-2-(4-cyano-phenylamino)-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy]-3,5-dimethyl-benzonitrile (28 mg, 0.060 mmol) in 1,2-dichlorobenzene (1 mL) was added aluminum chloride (40 mg, 0.30 mmol). The reaction mixture was stirred at 160 °C for 45 min and cooled to room temperature. The mixture was poured into ice water and extracted with CH₂Cl₂ (2 x 10 mL). The combined organic solution was washed with brine (10 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 50:50) yielded 6 mg (27%) of 4-[2-(4-cyano-phenylamino)-7H-pyrrolo[2,3-d]pyrimidin-4-yloxy]-3,5-dimethyl-benzonitrile as a tan solid.

25

20

Example 6

5

4-Cyclopropyl-2,6-dimethylphenol. To a suspension of (4-bromo-2,6-dimethylphenoxy) tert-butyldimethylsilane (668 mg, 2.12 mmol) and tetrakis(triphenylphosphine)palladium (122 mg, 0.11 mmol) in THF (20 mL) was added cyclopropyl zinc chloride (28.0 mL, 11.2 mmol). The mixture was heated at 80°C for 24 h and cooled to room temperature. The reaction was passed through a short pad of SiO₂ to remove the catalyst and the solution was concentrated to oil. The resulting oil was diluted in EtOAc (100 mL), washed with brine (100 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 90:10) yielded 370 mg (63%) of tert-butyl(4-cyclopropyl-2,6-dimethylphenoxy)dimethylsilane. To tert-butyl(4-cyclopropyl-2,6-

dimethylphenoxy)dimethylsilane (320 mg, 1.16 mmol) in THF (10 mL) was added a solution of tetrabutylammonium fluoride (5.0 mL, 1 M in THF, 5.0 mmol) and acetic acid (0.40 mL). The reaction was stirred at room temperature for 3 h and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 85:15) yielded 175 mg (93%) of 4-cyclopropyl-2,6-dimethylphenol as a light yellow oil.

2-chloro-6-(4-cyclopropyl-2,6-dimethylphenoxy)-9H-purine. To a solution of 4-cyclopropyl-2,6-dimethylphenol (263 mg, 1.62 mmol) in 1-methyl-2-pyrridone (3 mL) at 0°C was added NaH (65 mg, 1.62 mmol). The reaction mixture was stirred at room temperature for 30 min and a solution of 2,6-dichloropurine (102 mg, 0.54 mmol) in 1-methyl-2-pyrridone (2 mL) was added to the mixture. The mixture was heated at 100°C for 16 h and then cooled to room temperature. The reaction was poured into ice water and extracted with CHCl₃ (3 x 20 mL). The combined organic solution was washed with H₂O (20 mL) and brine (20 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (MeOH:CHCl₃ = 5:95) yielded 114 mg (67%) of 2-chloro-6-(4-cyclopropyl-2,6-dimethylphenoxy)-9H-purine.

10

5

4-(6-(4-cyclopropyl-2,6-dimethylphenoxy)-9H-purin-2-ylamino)benzonitrile. To a suspension of 2-chloro-6-(4-cyclopropyl-2,6-dimethylphenoxy)-9H-purine (28 mg, 0.088 mmol) and 4-aminobenzonitrile (42 mg, 0.35 mmol) in 2,2,2-trifluoroethanol (3 mL) in a sealed tube was added trifluoroacetic acid (0.056 mL, 0.70 mmol). The resulting solution was heated at 90°C for 3 days. The reaction was cooled to room temperature and concentrated to dryness. Silica gel chromatography (CH₂Cl₂:Acetone = 80:20) yielded 7 mg (20%) of 4-(6-(4-cyclopropyl-2,6-dimethylphenoxy)-9H-purin-2-ylamino)benzonitrile as a light yellow solid.

20

15

Example 7

5

10

15

20

2,4-dichloro-7H-pyrrolo[2,3-d]pyrimidine. To a suspension of 4-chloro-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (500 mg, 2.97 mmol) in 1,2-dichloroethane (40 mL) at -10°C under argon was added antimony chloride (750 mg, 3.29 mmol). After stirring for 5 min, tert-butylnitrite (2.50 mL, 20.8 mmol) was added to the solution. The reaction was stirred at -10°C for 3 h. The reaction was diluted with CHCl₃ (100 mL) and poured into ice water (50 mL). The CHCl₃ layer was separated, washed with brine (20 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 50:50) yielded 239 mg (43%) of 2,4-dichloro-7H-pyrrolo[2,3-d]pyrimidine as a tan solid.

2-chloro-4-(4-cyclopropyl-2,6-dimethylphenoxy)-7H-pyrrolo[2,3-d]pyrimidine. To a solution of 4-cyclopropyl-2,6-dimethylphenol (259 mg, 1.60 mmol) in THF (3 mL) at 0°C was added NaH (64 mg, 1.60 mmol). The reaction mixture was stirred at room temperature for 30 min and a solution of 2,4-dichloro-7H-pyrrolo[2,3-d]pyrimidine (100 mg, 0.53 mmol) in THF (2 mL) was added to the mixture. The mixture was heated at 80°C for 16 h and cooled to room temperature. The reaction was poured into ice water and extracted with CHCl₃ (3 x 20 mL). The combined organic solution was washed with H₂O (20 mL) and brine (20 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (Hexanes:EtOAc = 75:25) yielded 79 mg (48%) of 2-chloro-4-(4-cyclopropyl-2,6-dimethylphenoxy)-7H-pyrrolo[2,3-d]pyrimidine as a tan solid.

4-(4-(4-cyclopropyl-2,6-dimethylphenoxy)-7H-pyrrolo[2,3-d]pyrimidin-2-

ylamino)benzonitrile. To a suspension of 2-chloro-4-(4-cyclopropyl-2,6-dimethylphenoxy)-7H-pyrrolo[2,3-d]pyrimidine (75 mg, 0.24 mmol) and 4-aminobenzonitrile (113 mg, 0.96 mmol) in 2,2,2-trifluoroethanol (4 mL) in a sealed tube was added trifluoroacetic acid (0.15 mL, 1.92 mmol). The resulting solution was heated at 90°C for 3 days. The reaction was diluted with EtOAc (50 mL), washed with NaHCO₃ (20 mL) and brine (20 mL), dried with Na₂SO₄, and concentrated to dryness. Silica gel chromatography (CH₂Cl₂:Acetone = 90:10) yielded 15 mg (16%) of 4-(4-(4-cyclopropyl-2,6-dimethylphenoxy)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamino)benzonitrile as a tan solid.

Example 8

15

20

10

5

Scheme 1

Scheme 1 illustrate the synthesis of 9-deazaguanine by starting with commercially available 2-amino-6-methylpyrimidin-4(3*H*)-one and nitrated with nitric acid followed by treatment of the nitrated product with *N*,*N*-dimethylformamide dimethyl acetal (DMF-DMA) to afford the corresponding 2-(dimethylamino)methyleneimino derivative. It was then benzylated to produce 3-benzyl-2-[(dimethylamino)methyleneimino]-5-nitro-6-methylpyrimidin-4-one by treating with benzyl bromide and converted to benzylated-2,6-bis-dimethylaminomethylene derivative with DMF-DMA. Reductive cyclization with sodium

PCT/US2006/017677

hydrosulfite followed by de-protection with 3M NaOH and de-benzylation with Pd/C and NH₄CO₂H afforded 9-deazaguanine.

Scheme 2

5

10

15

Scheme 2 illustrates the 3 different pathways which provide the various substituted 9-deazapurines. Other products are synthesized by analogous methods, which a person skilled in the art could formulate, based on the reaction sequences given above. In certain cases, the person skilled in the art would see that protecting groups might be necessary. The synthetic scheme can be summarized as follows.

Benzylation of 9-deazaguanine followed by chlorination with $POCl_3$ gives the chlorinated 9-deazapurine product. This chlorinated intermediate can either be coupled with R2 (pathway 1) followed by diazotization with t-butyl nitrite; displaced with F; coupled with R3 and de-benzylated to give the product; or it can undergo pathway 2, which is diazotization

with t-butyl nitrite in the presence of antimony chloride followed by coupling with R2 and R3, followed by de-benzylation to afford the final product. Alternatively, pathway 3 provides for de-benzylation of the dichloro-9-deazapurine followed by the coupling with R2 and R3 respectively to provide the various substituted 9-deazapurine.

5

10

2-Amino-6-methyl-5-nitropyrimidin-4(3H)-one

To a mixture of 2-amino-6-methylpyrimidin-4(3H)-one (50 g, 0.4 mol) in 250 mL of H₂SO₄ at 0 °C was added 40 mL of HNO₃ with an additional funnel. After being stirred at room temperature for 3 h, the reaction mixture was slowly poured into 3.6 L of diethyl ether and stirred for 15 min. Decant the ether solution and added 1.0 L of ethyl acetate to the solid and stirred for 10 h. The solid (54.8 g, 81% yield) was filtered and used for next step without any further purification.

15

(E)-N,N-Dimethyl-N'-(4-methyl-5-nitro-6-oxo-1,6-dihydropyrimidin-2-methyl-N'-(4-methyl-5-nitro-6-oxo-1,6-dihydropyrimidin-2-methyl-N'-(4-methyl-5-nitro-6-oxo-1,6-dihydropyrimidin-2-methyl-1yl)formimidamide

To a suspension of 2-Amino-6-methyl-5-nitropyrimidin-4(3H)-one (54.8 g, 0.32 mol) in CH₂Cl₂ (461 mL) was added DMF-dimethylacetal (103.1 mL, 0.77 mol) and stirred at room temperature for 1.5 h. The reaction mixture was filtered, washed with CH₂Cl₂, and used for the next step without further purification (31.9 g, 44% yield).

20

(E)-N'(1-benzyl-4-methyl-5-nitro-6-oxo-1,6-dihydropyrimidin-2-yl)-N,Ndimethylformimidamide

To a suspension of (E)-N,N-Dimethyl-N'-(4-methyl-5-nitro-6-oxo-1,6-25 dihydropyrimidin-2-yl)formimidamide (53.4 g, 0.24 mmol) in DMF (690 mL) was added DBU (44.6 mL, 0.30 mol) and benzyl bromide (44.4 mL, 0.29 mol) and stirred at room

temperature for 1 h. The excess of DBU was neutralized with HCl, and the mixture was concentrated *in vacuo*. The residue was dissolved in methylene chloride and extracted twice with 2M HCl and water, then dried over Na₂SO₄ and concentrated. Trituration with ethanol afforded the crystalline product which was washed with ethanol to give the product (64.7 g, 86% yield) and used in the next step without further purification.

(E)-N'-(1-benzyl-4-((E)-2-(dimethylamino)vinyl)-5-nitro-6-oxo-1,6-dihydropyrimidin-2-yl)-N,N-dimethylformimidamide

To a solution of (E)-N'(1-benzyl-4-methyl-5-nitro-6-oxo-1,6-dihydropyrimidin-2-yl)
N,N-dimethylformimidamide (64.7 g, 0.2 mol) in DMF (254 mL) was added DMFdimethylacetal (54.5 mL, 0.41 mol). The reaction mixture was stirred for 3 h at 65 °C,
cooled, and the solvent was removed under reduced pressure. The residue was triturated with
ethanol, and the solid was collected by vacuum filtration (69.2 g, 91%) and used in the next
step without further purification.

15

20

5

(E)-N'-(3-benzyl-4-oxo-4,5-dihydro-3H-pyrrolo[3,2-d]pyrimidin-2-yl)-N,N-dimethylformimidamide

To a mixture of (E)-N'-(1-benzyl-4-((E)-2-(dimethylamino)vinyl)-5-nitro-6-oxo-1,6-dihydropyrimidin-2-yl)-N,N-dimethylformimidamide (43.0 g, 0.12 mol) in THF (151 mL) was added an aqueous saturated solution of $Na_2S_2O_4$ and stirred at room temperature overnight. Upon completion of the reaction, the solid was filtered and washed with THF to afford the product (21.2 g, 62% yield) which was used in the next step without further purification.

2-Amino-3-benzyl-3H-pyrrolo[3,2-d]pyrimidin-4(5H)-one

To a mixture of (E)-N'-(3-benzyl-4-oxo-4,5-dihydro-3*H*-pyrrolo[3,2-*d*]pyrimidin-2-yl)-*N*,*N*-dimethylformimidamide (21.2 g, 0.07 mol) in MeOH (382 mL) was added 3M NaOH (276 mL) and heated at 100 °C for 5 h. After completion of the reaction, the reaction mixture was cooled to 0 °C. The solid was filtered (15.8 g, 91%) and used in the next step without further purification.

2-Amino-3H-pyrrolo[3,2-d]pyrimidin-4(5H)-one

To a mixture of 2-amino-3-benzyl-3*H*-pyrrolo[3,2-*d*]pyrimidin-4(5*H*)-one (10 g, 0.04 mol) in MeOH (334 mL) was added 10% Pd/C (2 g), ammonium formate (13.2 g, 0.21 mmol) and heated at 75°C for 4 h. After completion of the reaction, the reaction mixture was cooled and filtered through a pad of Celite with hot 1:1 DMF/MeOH. The filtrate was concentrated *in vacuo* to provide the product as an off-white solid (6.2 g, 99%).

15

20

25

10

5

2-Amino-5-benzyl-3*H*-pyrrolo[3,2-*d*]pyrimidin-4(5*H*)-one

To a suspension of 2-amino-3*H*-pyrrolo[3,2-*d*]pyrimidin-4(5*H*)-one (336.7 mg, 2.0 mmol) in CH₂Cl₂ (14.3 mL) was added benzyl bromide (0.26 mL, 2.2 mmol) and TBABr (644 mg, 2.0 mmol). The reaction mixture was cooled to 0 °C, and to it was added 50% NaOH (1.7 mL). The resulting mixture was stirred for 2 h as it warmed from 0 °C to room temperature. Water was then added, and the solution was washed with CHCl₃. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by column chromatography, eluting with CH₂Cl₂/Acetone (5:1-1:1), afforded the product as a tan solid (423 mg, 82%)

5-benzyl-4-chloro-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-amine

A mixture of 2-amino-5-benzyl-3*H*-pyrrolo[3,2-*d*]pyrimidin-4(5*H*)-one (1.1 g, 7.4 mmol) and POCl₃ (7 mL, 74 mmol) was heated at 116 °C for 3 h. Upon completion of the reaction, the reaction mixture was poured into ice and extracted three times with ethyl acetate. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by column chromatography, eluting with CH₂Cl₂/Acetone (3:1), afforded the product as a white solid (490 mg, 40%).

10

15

20

5

5-benzyl-4-(mesityloxy)-5H-pyrrolo[3,2-d]pyrimidin-2-amine

To a stirred suspension of NaH (56 mg, 2.33 mmol) in dry NMP (2 mL) was added 2,4,6-trimethyl phenol (317 mg, 2.33 mmol). The mixture was stirred at room temperature for 30 min under argon. The reaction mixture was added to a solution of 5-benzyl-4-chloro-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-amine (200 mg, 0.78 mmol) in dry NMP (1.5 mL) and the resulting solution was heated at 90 °C for 16 h. After completion of the reaction, the reaction mixture was diluted with water and washed with EtOAc. The combined organic layers were washed with water, 2% NaOH, and brine and dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by column chromatography, eluting with hexanes/ethyl acetate (3:1) to give the product as a white solid (140 mg, 50%).

5-benzyl-2-fluoro-4-(mesityloxy)-5H-pyrrolo[3,2-d]pyrimidine

A solution of 5-benzyl-4-(mesityloxy)-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-amine (139.9 mg, 0.39 mmol) in pyridine (1.6 mL) was cooled to -50 °C and HF-pyr (8 mL) and *t*-butyl nitrite (0.19 mL, 1.56 mmol) was added dropwise. The reaction mixture was stirred at 50 °C to -30 °C for 1.5 h. Upon completion of the reaction, the reaction mixture was poured into K₂CO₃ (5 g), slowly added water and washed with CHCl₃ x 3. The combined organic layers were washed with brine, dried (Na₂SO₄), filtered and concentrated *in vacuo*. The crude product was purified by silica gel column chromatography, eluting with hexanes/ethyl acetate (2:1) to give the product as a white solid (116 mg, 82%).

10

15

20

25

4-(5-benzyl-4-(mesityloxy)-5H-pyrrolo[3,2-d]pyrimidin-2-ylamino)benzonitrile

A stirred suspension of NaH (63.8 mg, 2.66 mmol) in dry NMP (1.5 mL) was added 4-aminobenzylnitrile (188 mg, 2.66 mmol) and stirred at room temperature for 30 min under argon. The reaction mixture was added to a solution of 5-benzyl-2-fluoro-4-(mesityloxy)-5*H*-pyrrolo[3,2-*d*]pyrimidine (115 mg, 0.32 mmol) in dry NMP (1.7 mL) and stirred at room temperature for 2 h. After completion of the reaction, the resulting mixture was diluted with water and washed with EtOAc 3 times. The combined organic layers were washed with water, NH₄Cl, water x 2, and brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by column chromatography, eluting with 1% MeOH:CH₂Cl₂, which afforded the product as a tan solid (120 mg, 80%).

4-(4-(Mesityloxy)-5H-pyrrolo[3,2-d]pyrimidin-2-ylamino)benzonitrile

To a suspension of 4-(5-benzyl-4-(mesityloxy)-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-ylamino)benzonitrile (150 mg, 0.33 mmol) in 1,2-dichlorobenzene (13 mL) was added AlCl₃ (436 mg, 3.27 mmol). The reaction mixture was heated at 160 °C for 1.5 h during which the

reaction mixture became dark and homogeneous. Upon completion of the reaction, the reaction mixture was cooled and washed with NH₄Cl. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by column chromatography, eluting with Hexanes:Ethyl acetate (5:1-1:1) provided the product as a tan solid (27.8 mg, 23%).

Example 9

4-(2-Amino-5-benzyl-5H-pyrrolo[3,2-d]pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile

To a stirred suspension of NaH (155 mg, 6.47 mmol) in dry NMP (4 mL) was added 4-hydroxy-3,5-dimethylbenzonitrile (570 mg, 3.88 mmol), and the mixture was stirred at room temperature for 30 min under argon. The reaction mixture was added to a solution of 5-benzyl-4-chloro-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-amine (400 mg, 1.55 mmol) in dry NMP (4 mL) and heated at 160 °C for 16 h. After completion of the reaction, the resulting mixture was diluted with water and washed with EtOAc. The combined organic layers were washed with water, 2% NaOH, brine, dried over Na₂SO₄, filtered, and concentrated in vacuo. The crude product was purified by column chromatography, eluting with hexanes/ethyl acetate (2:1-1:4) to give the product as a light yellow solid (342 mg, 60%).

20

10

15

5

4-(5-benzyl-2-fluoro-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile

A solution of 4-(2-amino-5-benzyl-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile (319.4 mg, 0.87 mmol) in pyridine (3 mL) was cooled to -50 °C and

10

15

20

HF-pyr (15 mL) and t-butyl nitrite (0.42 mL, 3.46 mmol) were added dropwise. The reaction mixture was stirred at -50 °C to -20 °C for 1.5 h. Upon completion of the reaction, the mixture was poured into K₂CO₃ (8 g), diluted with water and washed with CHCl₃ x 3. The combined organic layers were washed with brine, dried (Na₂SO₄), filtered, and concentrated in vacuo. The crude product was purified by silica gel column chromatography, eluting with hexanes/ethyl acetate (2:1-1:1) which same the product as a light yellow solid (314 mg, 97%).

4-(5-benzyl-2-(4-cyanophenylamino)-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile

To a stirred suspension of NaH (101 mg, 4.21 mmol) in dry NMP (4 mL) was added 4-aminobenzylnitrile (299 mg, 2.53 mmol) and stirred at room temperature for 30 min under argon. The reaction mixture was added to a solution of 4-(5-benzyl-2-fluoro-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile (314 mg, 0.84 mmol) in dry NMP (4.4 mL) and stirred at room temperature for 2 h. After completion of the reaction, the resulting mixture was diluted with water and washed with EtOAc x 3. The combined organic layers were washed with water, NH₄Cl, water x 2, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by column chromatography, eluting with 1% MeOH:CH₂Cl₂, producing the product as a tan solid (320 mg, 80%).

4-(2-(4-Cyanophenylamino)-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile

To a suspension of 4-(5-benzyl-2-(4-cyanophenylamino)-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile (240 mg, 0.51 mmol) in 1,2-dichlorobenzene (20 mL) was added AlCl₃ (681 mg, 5.1 mmol). The reaction mixture was heated at 160 °C for 1.5 h, during which time the reaction mixture became dark and homogeneous. Upon completion of the reaction, the reaction mixture was cooled and washed with NH₄Cl. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated *in vacuo*. The crude product was purified by preparative TLC TLC eluting with Hexanes:Ethyl acetate (2.5:1) and produced the product as a pink solid (51 mg, 26%).

10 Example 10

5

15

20

$\hbox{5-benzyl-4-(mesitylthio)-5H-pyrrolo} \hbox{[3,2-d]} pyrimidin-2-amine$

To a stirred suspension of NaH (48 mg, 2 mmol) in dry NMP (2 mL) was added 2,4,6-trimethyl-benzene-1-thiol (191 mg, 1.2 mmol) The mixture was and stirred at room temperature for 30 min under argon. The reaction mixture was then added to a solution of 5-benzyl-4-chloro-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-amine (103 mg, 0.4 mmol) in dry NMP (2.5 mL) and heated at 60 °C for 16 h. After completion of the reaction, the resulting mixture was diluted with water and washed with EtOAc. The combined organic layers were washed with water, 2% NaOH, and brine; dried over Na₂SO₄; filtered; and concentrated in vacuo. The crude product was purified by column chromatography, eluting with hexanes/ethyl acetate (2:1-1:3) to give the product as a light yellow solid (131 mg, 88%).

5-benzyl-2-fluoro-4-(mesitylthio)-5H-pyrrolo[3,2-d]pyrimidine

A solution of 5-benzyl-4-(mesitylthio)-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-amine (131 mg, 0.35 mmol) in pyridine (1.6 mL) was cooled to -50 °C and added HF-pyr (8 mL) and *t*-butyl nitrite (0.17 mL, 1.4 mmol) dropwise. The reaction mixture was stirred at -50 °C to -40 °C for 1.5 h. Upon completion of the reaction, the reaction was poured into K₂CO₃ (5 g), slowly added water and washed with CHCl₃ x 3. The combined organic layers were washed with brine, dried (Na₂SO₄), filtered and concentrated *in vacuo*. The crude product was purified by silica gel column chromatography, eluting with hexanes/ethyl acetate (5:1-1:1) to give the product as an off-white solid (94 mg, 71%).

10

15


20

25

5

4-(5-benzyl-4-(mesitylthio)-5H-pyrrolo[3,2-d]pyrimidin-2-ylamino)benzonitrile

To a stirred suspension of NaH (30 mg, 1.25 mmol) in dry NMP (1.5 mL) was added 4-aminobenzylnitrile (87.4 mg, 0.74 mmol) and stirred at room temperature for 30 min under argon. The reaction mixture was added to a solution of 5-benzyl-2-fluoro-4-(mesitylthio)-5*H*-pyrrolo[3,2-*d*]pyrimidine (93 mg, 0.25 mmol) in dry NMP (1 mL) and stirred at room temperature for 2 h. After completion of the reaction, the resulting mixture was diluted with water and washed with EtOAc x 3. The combined organic layers were washed with water, NH₄Cl, water x 2, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by preparative TLC TLC, eluting with Hexanes:Ethyl acetate (1.5:1) afforded the product as a tan solid (12.6 mg, 11%).



4-(4-(mesitylthio)-5H-pyrrolo[3,2-d]pyrimidin-2-ylamino)benzonitrile

To a suspension of 4-(5-benzyl-4-(mesitylthio)-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-ylamino)benzonitrile (9.2 mg, 0.03 mmol) in 1,2-dichlorobenzene (1 mL) was added AlCl₃

5

10

15

20

(26 mg, 0.3 mmol). The reaction mixture was heated at 160 °C for 1.5 h, which the reaction mixture became dark and homogeneous. Upon completion of the reaction, the reaction mixture was cooled and washed with NH₄Cl. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by preparative TLC TLC, eluting with Hexanes:Ethyl acetate (2.5:1) produced the product as a pink solid (7.7 mg, 20%).

Example 11

5-benzyl-2,4-dichloro-5*H*-pyrrolo[3,2-*d*]pyrimidine

To a suspension of 5-benzyl-4-chloro-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-amine (641 mg, 2.5 mmol) in 1,2-dichloroethane (35 mL) was cooled to -10 °C SbCl₃ (850 mg, 3.7 mmol) was added. The reaction mixture was stirred for 5 min. *t*-butyl nitrite (2.1 mL, 17.4 mmol) was added dropwise and the stirred mixture was from -10 °C to room temperature for 5 h. Upon completion of the reaction, the reaction mixture was poured into ice water and washed with CH₂Cl₂. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated *in vacuo*. The crude product was purified by silica gel column chromatography eluting with Hexanes:Ethyl acetate (9:1-1:1), and gave the product as an off-white solid (528 mg, 77%).

2,4-Dichloro-5H-pyrrolo[3,2-d]pyrimidine

To a suspension of 5-benzyl-2,4-dichloro-5*H*-pyrrolo[3,2-*d*]pyrimidine (177 mg, 0.64 mmol) in 1,2-dichlorobenzene (20 mL) was added AlCl₃ (852 mg, 6.4mmol). The reaction mixture was heated at 160 °C for 1.5 h, during which the reaction mixture became dark and homogeneous. Upon completion of the reaction, the reaction mixture was cooled, added CHCl₃ and washed with NH₄Cl. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Added Hexanes and filtered off the

product as purple solids (100 mg, 80%) and used for the next step without further purification.

2-Chloro-4-(4-cyclopropyl-2,6-dimethylphenoxy)-5*H*-pyrrolo[3,2-*d*]pyrimidine

To a stirred suspension of NaH (25 mg, 0.64 mmol) in dry NMP (1.5 mL) was added 4-cyclopropyl-2,6-dimethylphenol (103 mg, 0.64 mmol) and the resolution mixture was stirred at room temperature for 30 min under argon. The reaction mixture was added to a solution of 2,4-dichloro-5*H*-pyrrolo[3,2-*d*]pyrimidine (120 mg, 0.64 mmol) in dry NMP (1.7 mL) and heated at 90 °C for 16 h. After completion of the reaction, the resulting mixture was diluted with water and washed with EtOAc. The combined organic layers were washed with water, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by column chromatography, eluting with hexanes/ethyl acetate (4:1-2:1), to give the product as a light yellow solid (20.2 mg, 8%).

15

20

10

5

4-(4-(4-Cyclopropyl-2,6-dimethylphenoxy)-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-ylamino)benzonitrile

In a sealed tube was placed 2-chloro-4-(4-cyclopropyl-2,6-dimethylphenoxy)-5*H*-pyrrolo[3,2-*d*]pyrimidine (20 mg, 0.064 mmol), 4-aminobenzonitrile (31 mg, 0.26 mmol), TFE (0.21 mL) and TFA (0.04 mL, 0.51 mmol). The reaction mixture was stirred at 90 °C for 16 h. Upon completion of the reaction, the resulting mixture was diluted with water and washed with EtOAc. The combined organic layers were washed with NaHCO₃, brine, dried

over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by preparative TLC TLC, eluting with 5 % Acetone/CH₂Cl₂ to give the product as a light yellow solid (10.5 mg, 45%).

5 Example 12

10

15

2-Chloro-4-(mesityloxy)-5-methyl-5H-pyrrolo[3,2-d]pyrimidine

To a stirred suspension of NaH (8.9 mg, 0.22 mmol) in dry NMP (1.0 mL) was added 2,4,6-trimethyl phenol (30.2 mg, 0.22 mmol) and stirred at room temperature for 30 min under argon. The reaction mixture was added to a solution of 2,4-dichloro-5-methyl-5*H*-pyrrolo[3,2-*d*]pyrimidine (44.6 mg, 0.22 mmol) in dry NMP (1.0 mL) and heated at 90 °C for 16 h. After completion of the reaction, the resulting mixture was cooled, diluted with water and washed with EtOAc. The combined organic layers were washed with water, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by column chromatography, eluting with hexanes/ethyl acetate (5:1-2:1), to give the product as a light yellow solid (52.7 mg, 80%).

4-(4-(Mesityloxy)-5-methyl-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-ylamino)benzonitrile

In a sealed tube was added 2-chloro-4-(mesityloxy)-5-methyl-5*H*-pyrrolo[3,2-*d*]pyrimidine (52.7 mg, 0.18 mmol), 4-aminobenzonitrile (83 mg, 0.70 mmol), TFE (1.0 mL) and TFA (0.11 mL, 1.44 mmol). The reaction mixture was stirred at 90 °C for 48 h. Upon completion of the reaction, the resulting mixture was cooled, diluted with water and washed

with EtOAc. The combined organic layers were washed with NaHCO₃, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by preparative TLC, eluting with hexanes:ethyl acetate (5:1-2:1), to give the product as a light yellow solid (65.7 mg, 95%).

5


10

15

20

4-(2-Chloro-5-methyl-5H-pyrrolo[3,2-d]pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile

To a stirred solution of NaH (42.1 mg, 1.05 mmol) in dry NMP (2.5 mL) was added 4-hydroxy-3,5-dimethylbenzonitrile (154.7 mg, 1.05 mmol) and stirred at room temperature for 30 min under argon. The reaction mixture was added to a solution of 2,4-dichloro-5-methyl-5*H*-pyrrolo[3,2-*d*]pyrimidine (211.3 mg, 1.05 mmol) in dry NMP (2.7 mL) and heated at 160 °C for 16 h. After completion of the reaction, the resulting mixture was diluted with water and washed with EtOAc. The combined organic layers were washed with water, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by column chromatography, eluting with hexanes/ethyl acetate (3:1-1:1), to give the product as a light yellow solid (294 mg, 85%).



 $\label{eq:continuous} \mbox{4-(2-(4-Cyanophenylamino)-5-methyl-5$$H$-pyrrolo[3,2-$d]$ pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile$

In a sealed tube was added 4-(2-chloro-5-methyl-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile (294 mg, 0.94 mmol), 4-aminobenzonitrile (455 mg, 3.77 mmol), TFE (3.1 mL) and TFA (0.58 mL, 7.52 mmol). The reaction mixture was stirred at 90 °C for 48 h. Upon completion of the reaction, the resulting mixture was cooled, diluted

with water and washed with EtOAc. The combined organic layers were washed with NaHCO₃, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by preparative TLC, eluting with hexanes:ethyl acetate (4:1-1:2), to give the product as an off-white solid (133 mg, 40%).

5

Example 13

10

15

20

1-(4-(5-benzyl-2-chloro-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-yloxy)-3,5-dimethylphenyl)ethanone

To a stirred solution of NaH (31 mg, 0.78 mmol) in dry NMP (2 mL) was added 1-(4-hydroxy-3,5-dimethylphenyl)ethanone (127 mg, 0.78 mmol) and stirred at room temperature for 30 min under argon. The reaction mixture was added to a solution of 5-benzyl-2,4-dichloro-5*H*-pyrrolo[3,2-*d*]pyrimidine (216 mg, 0.78 mmol) in dry NMP (2.4 mL) and heated at 160 °C for 16 h. After completion of the reaction, the resulting mixture was diluted with water and washed with EtOAc. The combined organic layers were washed with water, 2% NaOH, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by column chromatography, eluting with hexanes/ethyl acetate (4:1-2:1), to give the product as a light yellow solid (111 mg, 35%).

5

10

4-(4-(4-Acetyl-2,6-dimethylphenoxy)-5-benzyl-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-ylamino)benzonitrile

In a sealed tube was added 1-(4-(5-benzyl-2-chloro-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-yloxy)-3,5-dimethylphenyl)ethanone (111 mg, 0.27 mmol), 4-aminobenzonitrile (129 mg, 1.1 mmol), TFE (1.7 mL) and TFA (0.2 mL, 2.16 mmol). The reaction mixture was stirred at 90 °C for 16 h. Upon completion of the reaction, the resulting mixture was cooled, diluted with water, and washed with EtOAc. The combined organic layers were washed with NaHCO₃ and brine; dried over Na₂SO₄; filtered; and concentrated *in vacuo*. The crude product was purified by silica gel column chromatography, eluting with hexanes:ethyl acetate (9:1-100 % EtOAc), to give the product as an off-white solid (68 mg, 51%).

4-(4-(4-Acetyl-2,6-dimethylphenoxy)-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-ylamino)benzonitrile

To a suspension of 4-(4-(4-acetyl-2,6-dimethylphenoxy)-5-benzyl-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-ylamino)benzonitrile (65 mg, 0.13 mmol) in 1,2-dichlorobenzene (5.3 mL) was added AlCl₃ (178 mg, 1.3 mmol). The reaction mixture was heated at 160 °C for 1.5 h, after which time the reaction mixture became dark and homogeneous. Upon completion of the reaction, the reaction mixture was cooled, CHCl₃ was added, and the mixture was washed with NH₄Cl. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel column chromatography, eluting with hexanes:ethyl acetate (3:1), to give the product as a brown solid (41 mg, 77%).

Example 14

25

15

20

$\label{eq:continuous} \begin{tabular}{l} 1-(4-(5-benzyl-2-5H-pyrrolo[3,2-d]pyrimidin-4-yloxy)-3,5-dimethylphenyl)-N,N-dimethylmethanamine \end{tabular}$

To a stirred solution of NaH (80.4 mg, 1.0 mmol) in dry NMP (3 mL) was added 4- ((dimethylamino)methyl)-2,6-dimethylphenol (216.4 mg, 1.0 mmol) and the mixture was stirred at room temperature for 30 min under argon. The reaction mixture was added to a solution of 5-benzyl-2,4-dichloro-5*H*-pyrrolo[3,2-*d*]pyrimidine (216 mg, 0.78 mmol) in dry NMP (2.6 mL) and heated at 120 °C for 16 h. After completion of the reaction, the resulting mixture was diluted with water and washed with EtOAc. The combined organic layers were washed twice with water, washed with, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel column chromatography, eluting with MeOH/CH₂Cl₂ (10%-30%), to give the product as a tan solid (71 mg, 17%).

15

20

5

10

4-(5-benzyl-4-(4-((dimethylamino)methyl)-2,6-dimethylphenoxy)-5H-pyrrolo[3,2-d]pyrimidin-2-ylamino)benzonitrile

In a sealed tube was added 1-(4-(5-benzyl-2-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-yloxy)-3,5-dimethylphenyl)-*N*,*N*-dimethylmethanamine (70.7 mg, 0.17 mmol), 4-aminobenzonitrile (78.9 mg, 0.67 mmol), TFE (1.1 mL) and TFA (0.1 mL, 1.3 mmol). The reaction mixture was stirred at 90 °C for 16 h. Upon completion of the reaction, the resulting mixture was cooled, diluted with water, and washed with EtOAc. The combined organic layers were washed with NaHCO₃ solution and with brine, dried over Na₂SO₄, filtered, and concentrated

in vacuo. The crude product was purified by silica gel column chromatography, eluting with MeOH/CH₂Cl₂ (20%-40%), to give the product as a tan solid (17 mg, 20%).

5 4-(4-(4-((dimethylamino)methyl)-2,6-dimethylphenoxy)-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-ylamino)benzonitrile

The benzyl group was removed according to the same procedure as described for example 13.

10

15

20

Example 15

5-benzyl-2-chloro-4-(2,6-dimethyl-4-nitrophenoxy)-5H-pyrrolo[3,2-d] pyrimidine

To a stirred solution of NaH (61.9 mg, 2.6 mmol) in dry NMP (4.7 mL) was added 2,6-dimethyl-4-nitrophenol (258.9 mg, 1.55 mmol) and stirred at room temperature for 30 min under argon. The reaction mixture was added to a solution of 5-benzyl-2,4-dichloro-5*H*-pyrrolo[3,2-*d*]pyrimidine (431 mg, 1.55 mmol) in dry NMP (4 mL) and heated at 90 °C for 16 h. After completion of the reaction, the resulting mixture was diluted with water and washed with EtOAc. The combined organic layers were washed twice with water, washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel column chromatography, eluting with hexanes:ethyl acetate (3:1-1:1), to give the product as a white solid (598 mg, 94%).

4-(5-benzyl-4-(2,6-dimethyl-4-nitrophenoxy)-5H-pyrrolo[3,2-d]pyrimidin-2-ylamino)benzonitrile

In a sealed tube was added 5-benzyl-2-chloro-4-(2,6-dimethyl-4-nitrophenoxy)-5*H*-pyrrolo[3,2-*d*]pyrimidine (598 mg, 1.46 mmol), 4-aminobenzonitrile (691 mg, 5.85 mmol), TFE (9.1 mL) and TFA (1.97 mL, 11.7 mmol). The reaction mixture was stirred at 90 °C for 16 h. Upon completion of the reaction, the resulting mixture was cooled, diluted with water and washed with EtOAc. The combined organic layers were washed with NaHCO₃, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel column chromatography, eluting with hexanes:ethyl acetate (3:1-1:1), to give the product (442 mg, 60%).

Example 16

15

10

5

$\label{eq:continuous} \mbox{4-(4-(4-Acetyl-2,6-dimethylphenoxy)-7-chloro-5$$H$-pyrrolo[3,2-$d]$ pyrimidin-2-ylamino) benzonitrile$

To a solution of 4-(4-(4-acetyl-2,6-dimethylphenoxy)-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-ylamino)benzonitrile (13 mg, 0.03 mmol) in CH₂Cl₂ (1 mL) was added NCS (4.4 mg, 0.03 mmol) and the mixture refluxed for 16 h. After the completion of the reaction, the solvent was concentrated and purified by preparative TLC eluting with hexanes:ethyl acetate (2:1) to give the product (4.2 mg, 30%).

Example 17

5 4-(7-Chloro-4-(mesitylthio)-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-ylamino)benzonitrile

To a solution of 4-(4-(mesitylthio)-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-ylamino)benzonitrile (10 mg, 0.02 mmol) in CH₂Cl₂ (5 mL) was added NCS (2.8 mg, 0.02 mmol) and the resolution mixture refluxed for 16 h. After the completion of the reaction, the solvent was concentrated and purified by preparative TLC, eluting with hexanes:ethyl acetate (3:1), to give the product (8.8 mg, 88%).

Example 18

15

20

10

$4\hbox{-}(7\hbox{-}bromo\hbox{-}4\hbox{-}(mesitylthio)\hbox{-}5H\hbox{-}pyrrolo[3,2\hbox{-}d] pyrimidin\hbox{-}2\hbox{-}ylamino) benzonitrile}$

To a solution of 4-(4-(mesitylthio)-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-ylamino)benzonitrile (22.7 mg, 0.06 mmol) in CH₂Cl₂(10 mL) was added NBS (10.5 mg, 0.06 mmol) and the resultant mixture was refluxed for 16 h. After completion of the reaction, the solvent was concentrated and purified by reversed phase HPLC to give the product as a white solid (6.4 mg, 23%).

Example 19

$\textbf{4-} (7-\text{chloro-4-} (\text{mesityloxy}) \textbf{-5-methyl-5} \textbf{\textit{H-pyrrolo}} [\textbf{3,2-d}] \textbf{pyrimidin-2-ylamino}) \textbf{benzonitrile}$

To a solution of 4-(4-(mesityloxy)-5-methyl-5*H*-pyrrolo[3,2-*d*]pyrimidin-2-ylamino)benzonitrile (17.3 mg, 0.05 mmol) in CH₂Cl₂ (5 mL) was added NCS (6.03 mg, 0.05 mmol) and the resultant mixture was refluxed for 16 h. After completion of the reaction, the solvent was concentrated and purified by preparative TLC, eluting with hexanes:ethyl acetate (3:1), to give the product as an off-white solid (3.4 mg, 6%).

10 Example 20

15

20

4-(7-Chloro-2-(4-cyanophenylamino)-5-methyl-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile

To a solution of 4-(2-(4-cyanophenylamino)-5-methyl-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile (21.5 mg, 0.06 mmol) in CH₂Cl₂ (3 mL) was added NCS (7.3 mg, 0.06 mmol) and the resultant mixture was refluxed for 16 h. After completion of the reaction, the solvent was concentrated and purified by preparative TLC, eluting with Hexanes:Ethyl acetate (3:1), to give the product as a light yellow solid (13.2 mg, 56%).

Example 21

$4-(7-{\bf Bromo-2-(4-cyanophenylamino)-5-methyl-5} \\ H-pyrrolo[3,2-d] pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile$

To a solution of 4-(2-(4-cyanophenylamino)-5-methyl-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile (58 mg, 0.15 mmol) in CH₂Cl₂ (8 mL) was added NBS (29 mg, 0.16 mmol) and the resultant mixture was refluxed for 16 h. After completion of the reaction, the solvent was concentrated and purified by preparative TLC, eluting with hexanes:ethyl acetate (2:1), to give the product as a yellow solid (40 mg, 57%).

10

5

Example 22

$\label{eq:continuous} \mbox{4-(7-Chloro-2-(4-cyanophenylamino)-5H-pyrrolo[3,2-d]pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile$

To a solution of 4-(2-(4-cyanophenylamino)-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-yloxy)-3,5-dimethylbenzonitrile (28.1 mg, 0.07 mmol) in CH₂Cl₂ (4 mL) was added NCS (9.9 mg, 0.07 mmol) and the resultant mixture was refluxed for 16 h. After completion of the reaction, the solvent was concentrated and purified by preparative TLC eluting with hexanes:ethyl acetate (2:1), to give the product as a pink solid (20 mg, 65%).

15

Biological Activity

Inhibition of HIV-1 Reverse Transcriptase

5

10

15

Numerous compounds were screened for inhibitory activity against human immunodeficiency virus type 1 (HIV-1) using a high throughput cell-based assay using HIV-1 expressing firefly luciferase as a reporter gene and pseudotyped with vesicular stomatitis virus envelope glycoprotein (VSV-G). Experimental procedures were essentially as described by Connor *et al.* in *Journal of Virology* (1996), 70: 5306-5311 (Characterization of the functional properties of env genes from long-term survivors of human immunodeficiency virus type 1 infection), and Popik *et al.* in *Journal of Virology* (2002), 76: 4709-4722 (Human immunodeficiency virus type 1 uses lipid raft-co-localized CD4 and chemokine receptors for productive entry into CD4+ T cells). It should be particularly appreciated that the virus contains two introduced mutations in the RT gene (K103N and Y181C, created by PCR mutagenesis) that render the virus highly resistant to current non-nucleoside HIV-1 drugs. Virus stocks were generated by cotransfection of plasmid DNA encoding VSV-G with vector pNL4-3Env(-)Luc(+) into 293T cells. Sixty-four hours after transfection, virus-containing medium was collected by centrifugation and stored frozen at -80° C.

20

25

30

35

HeLa cells were infected with the VSV-G pseudotyped virus in the presence of screening compounds in a 384-well microtiter plate format. Forty-eight hours after initial infection, lysis buffer and Luciferase Assay Reagent (Promega) was added to the cells and luciferase activity was determined by counting the resultant luminescence using a LJL luminometer. Since the luciferase gene is carried in the virus genome, its expression level directly reflects the virus replication level in the presence of a compound.

To evaluate the activity of the compounds against wild type HIV-1, the HeLa-JC53 cell line that expresses high levels of CD4 and CCR5 (see e.g., Platt *et al.* in *Journal of Virology* (1998), 72: 2855-2864: Effect of CCR5 and CD4 cell surface concentrations on infection by macrophagetropic isolates of human immunodeficiency virus type 1) was modified by isolation of a stable cell line that expresses luciferase under the control of the HIV-1 promoter (long terminal repeat, i.e., LTR). HIV-1 infection of this cell line stimulates the transcription of luciferase from the HIV-1 promoter and the luciferase gene expression level is proportional to the level of virus replication (Harrington *et al.* in *Journal of Virology*

WO 2006/122003 PCT/US2006/017677 **5**6

Methods (2000), 88: 111-115: Direct detection of infection of HIV-1 in blood using a centrifugation-indicator cell assay; and Roos et al. in Virology (2000), 273: 307-315: LuSIV cells: a reporter cell line for the detection and quantitation of a single cycle of HIV and SIV replication). Procedures for virus infection, compound testing and luciferase activity determination were the same as for the VSV-G pseudotyped HIV-1.

Two approaches were used to evaluate the cytotoxicity of the positive compounds discovered in the HIV-1 virus assays. The first approach employed another modified HeLa-JC53 cell line that constitutively expresses high level of luciferase without virus infection. The level of luciferase expression in these cells served as an indicator for cell replication in the presence of the compounds. Procedures for compound testing and luciferase activity determination were the same as for the virus infection tests. The other toxicity assay utilized HeLe-JC53 cells and a commercially available MTS assay kit (Promega) that measures the mitochondria function of the cells.

Results

5

10

15

20

The results are listed in Table A as EC50 (nM) and IC50 (nM). Table legend: A is < 10, B is between 10 and 100, C is > 100, ND is not determined. Note that many compounds of this invention exhibit activities on wild-type (WT) and resistant mutants below 10 nM.

Table A

Cpd	Structure	EC ₅₀ WT (nM)	EC ₅₀ Y181C (nM)	EC ₅₀ Y188L (nM)	EC ₅₀ L100I- K103N (nM)
1	CLogP: 6.31119	A	В	В	В
2	CLogP: 7.1942	A	В	В	С
3	NH CN CN CLogP: 5.24519	A	A	A	A

Cpd	Structure	EC ₅₀ WT (nM)	EC ₅₀ Y181C (nM)	EC ₅₀ Y188L (nM)	EC ₅₀ L100I- K103N (nM)
4	CLogP: 7.00339	A	В	В	В
5	CLogP: 5.71888	A	A	A	A
6	NH CN	A	В	В	В
7	N-NH N CN CLogP: 5.85982	A	A	В	В
8	CN CN CN CN CN CCN CCN CCN CCN CCN CCN	В	С	С	С
9	CLogP: 5.81219	A	С	С	С
10	CLogP: 6.65919	A	В	В	В
11	CI CLogP: 6.5252	A	В	A	В
12	CLogP: 4.43498	В	С	С	С

		EC ₅₀	EC ₅₀	EC ₅₀	EC ₅₀
Cpd	Structure	WT (nM)	Y181C (nM)	Y188L (nM)	L100I- K103N (nM)
		(11141)		,	KIOSIA (IIIAI)
13	CN CLogP: 5.18408	A	A	A	A
14	OMe CLogP: 6.24839	A	С	С	С
15	NH NH ₂ CLogP: 4.67978	В	С	С	С
16	CLogP: 6.01359	С	С	С	С
17	N-NH ON CLogP: 4.79382	A	В	A	В
18	CLogP: 7.04473	В	С	С	С
19	CLogP: 6.16288	A	A	A	В
20	NH N CN HN N N N N N N N N N N N N N N N N N N	A	В	С	С
21	CI CLogP: 6.10388	A	В	В	С

Cpd	Structure	EC ₅₀ WT (nM)	EC ₅₀ Y181C (nM)	EC ₅₀ Y188L (nM)	EC ₅₀ L100I- K103N (nM)
22	CLogP: 7.12035	· A	С	С	С
23	CLogP: 6.50038	A	A	A	В
24	N	В	С	C	С
25	NH CN CN CLogP: 4.6528B	A	A	A	В
26	CN CLogP: 5.71308	A	В	В	С
27	NH N CN CN N N N N N N N N N N N N N N N	A	В	В	
28	NH CN CN HN N N CN CLogP; 6.94438	A	A	В	В
29	CLogF: 6.01638	A	С	С	С
30	NH N CN CN CLogP: 5.42063	С	С	С	С

Cpd	Structure	EC ₅₀ WT (nM)	EC ₅₀ Y181C (nM)	EC ₅₀ Y188L (nM)	EC ₅₀ L100I- K103N (nM)
31	CLogP: 6.34283	A	В	A	В
32	CLogP: 6.67638	A	В	· C	С
33	CLogP: 6,75519	A	В	В	В
34	NH F CLogP: 6.41174	A	A	В	В
35	OMe CLogP: 6.18265	A	В	С	С
36	NH N CN CLogP: 5.71688	A	В	С	С
37	CLogP: 6.52119	A	A	A	A
38	NH CF ₃ CF ₃ CLogP: 7.16147	В	С	С	С
39	NH ONN CF ₃ Clo ₀ P; 7.16147	С	С	С	С

Cpd	Structure	EC ₅₀ WT (nM)	EC ₅₀ Y181C (nM)	EC ₅₀ Y188L (nM)	EC ₅₀ L100I- K103N (nM)
40	NH CN CF ₃ CLogP: 6.60082	В	В	С	В
41	CN CLogP: 7.16208	Α	A	В	В
42	CLogP: 5.45519	A	A	A	A
43	CLogP: 0.95418	В	В	С	В
44	OLogP: 7.19876	В	С	C	С
45	CLogP: 5.46143	A	A	A	В
46	CN CLogP: 6.29507	С	С	C	С
47	CLogP: 8.67208	В	В	С	С

Cpd	Structure	EC ₅₀ WT (nM)	EC ₅₀ Y181C (nM)	EC ₅₀ Y188L (n M)	EC ₅₀ L100I- K103N (nM)
48	CLogP: 9.11123	В	В	С	С
49	CN CLogP: 5.18373	A	A	A	A
50	MeO OMe CLogF: 4.02888	, A	A	В	С
51	HN N CN S N N CLogP: 7.4042	A	A	A	В.
52	HN N CN CN CN CLogP: 6.96519	A	A	A	A
53	CI N CN S N N Y	A	В	В	С
54	CLogP: 9.83084	A	В	В	С
55	Br N CN CLogP: 8.28777	A	В	В	С

	Structure	EC ₅₀ WT	EC ₅₀ Y181C (nM)	EC ₅₀ Y188L (n M)	EC ₅₀ L100I-
Cpd		(nM)			K103N (nM)
56	CLogP: 7.88168	В	В	В	В
57	CLogP: 6.46008	A	A	A	A
58	CLogP: 8.59208	В	В	C	С
59	CI ONN H CN CLcgP: 7.17968	В	В	В	В
60	CLogP: 7.16808	A	A	В	В
61	CLogP: 5.46119	A	A	A	A
62	CLogP: 7.39368	A	A	A	A
63	Cl ClogP: 8.44208	В	В	С	С

Cpd	Structure	EC ₅₀ WT (nM)	EC ₅₀ Y181C (nM)	EC ₅₀ Y188L (nM)	EC ₅₀ L100I- K103N (nM)
64	CLogP: 8 22808	A	В	С	С
65	CLogP: 6.02219	A	A	A	С
66	HN CN CN CN CLogP: 6.73519		A	A	A
67	Cl N CLcgP: 6.19473	A	A	В	A
68	ON H CLogP: 5.39408	A	A	A	A
69	CN CLogP: 6.11368	A	A	A	A
70	CLogP: 6.26368	A	A	A	A
71	CLogP: 6.36919	A	A	A	A

Contemplated Compounds and Prophetic Examples

In addition to the examples listed above, this invention provides or contemplates many compounds, examples of which are shown in the tables that follow.

5

Table 1 Contemplated Compounds of Formula IA-1

10

IA-1

	Ar la	V	W	Z
1.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	Н	CH ₃
2.	o, o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	benzyl	CH ₃
3.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	benzyl	H
4.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	3-Me-benzyl	· CH ₃
5.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	4-Me-benzyl	H
6.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	3-MeO-benzyl	H
7.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	4-MeO-benzyl	CH ₃
8.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	H	H
9.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	H	Br
10.	o, o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	cyclopropyl	CH ₂ CH ₃
11.	o, o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	CH ₂ CF ₃	CH₂CH₃
12.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	H	\mathbf{CH}_3
13.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	benzyl	CH₃
14.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	benzyl	H
15.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	3-Me-benzyl	cyclopropyl
16.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	3-MeO-benzyl	benzyl
17.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	H	H
18.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	C≡CCH ₃	CH ₂ CH ₃	$\mathrm{CH_3}$
19.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	Cl	CH ₂ CH=CH ₂	H
20.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ CH ₃	CH ₂ CH=CH ₂	H
21.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	Cl	CH ₂ CH ₃	CH ₂ CH ₃
22.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	C1	H	Н
23.	4-cyclopropylnaphth-l-yl	CN	H	CH ₃
24.	4-cyclopropylnaphth-l-yl	CN	benzyl	CH_3
25.	4-cyclopropylnaphth-l-yl	CN	benzyl	Н
26.	4-cyclopropylnaphth-l-yl	CN	H	H
27.	4-cyclopropylnaphth-l-yl	CH=CHCN	H	CH ₃
28.	4-cyclopropylnaphth-l-yl	CH=CHCN	benzyl	CH ₃
29.	4-cyclopropylnaphth-l-yl	CH=CHCN	benzyl	H
30.	4-cyclopropylnaphth-l-yl	CH=CHCN	H	H
31.	4-cyclopropylnaphth-l-yl	SO ₂ NHCH ₃	CH ₂ CN	F

	Ar	V	W	Z
32.	4-cyclopropylnaphth-l-yl	SO ₂ NHCH ₃	cyclopropyl	Cl
33.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₂ CH ₂ CN	Br
34.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₂ CN	benzyl
35.	o,o'-di-CH ₃ O-p-CN-phenyl	C≡CCH ₃	3-MeO-benzyl	F
36.	o,o'-di-CH ₃ O-p-CN-phenyl	F	3-Me-benzyl	Cl
37.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	H	CH ₃
38.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	benzyl	CH ₃
39.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	benzyl	H
40.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	Н	H
41.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	H	$\mathrm{CH_{3}}$
42.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	benzyl	CH_3
43.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	benzyl	H
44.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	H	H
45.	o,o'-di-CH ₃ -p-CN-phenyl	CN	H	CH ₃
46.	o,o'-di-CH ₃ -p-CN-phenyl	CN	benzyl	CH ₃
	·		3,5-di MeO-	
47.	o,o'-di-CH ₃ -p-CN-phenyl	CN	benzyl	CH ₃
48.	o,o'-di-CH ₃ -p-CN-phenyl	CN	benzyl	H
49.	o,o'-di-CH ₃ -p-CN-phenyl	CN	H	H
50.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	H	CH ₃
51.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	benzyl	CH ₃
52.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	benzyl	H
53.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	H	H
54.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	H	F
55.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	benzyl	F
56.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	benzyl	F
57.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	H	F
58.	4-cyclopropylnaphth-l-yl	CN	H	F
59.	4-cyclopropylnaphth-l-yl	CN	benzyl	F
60.	4-cyclopropylnaphth-l-yl	CH=CHCN	H	F
61.	4-cyclopropylnaphth-l-yl	CH=CHCN	benzyl	F
62.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	H	F
63.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	benzyl	F
64.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	H	F
65.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	benzyl	F
66.	o,o'-di-CH ₃ -p-CN-phenyl	CN	H	F
67.	o,o'-di-CH ₃ -p-CN-phenyl	CN	benzyl	F
68.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	H	F F
69.	o, o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	benzyl	120
70.	o, o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	H	CH ₃
71.	o, o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	benzyl	CH ₃
72.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	benzyl H	<u>Н</u> Н
73.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	H	
74.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂		CH ₃
75.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F F	benzyl	CH ₃
76.	o, o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	benzyl	H
77.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	J F	H	H

	Ar	V	W	Z
78.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	H	CH ₃
79.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	benzyl	CH ₃
80.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	benzyl	H
81.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	H	H
82.	4-cyclopropylnaphth-l-yl	F	H	CH ₃
83.	4-cyclopropylnaphth-l-yl	F	benzyl	CH ₃
84.	4-cyclopropylnaphth-l-yl	F	benzyl	H
85.	4-cyclopropylnaphth-l-yl	F	H	H
86.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	H	CH ₃
87.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	benzyl	CH ₃
88.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	benzyl	H
89.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	H	Н
90.	o,o'-di-CH ₃ O-p-CN-phenyl	F	H	CH ₃
91.	o,o'-di-CH ₃ O-p-CN-phenyl	F	benzyl	CH ₃
92.	o,o'-di-CH ₃ O-p-CN-phenyl	F	benzyl	Н
93.	o,o'-di-CH ₃ O-p-CN-phenyl	F	H '	H
94.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	H	CH ₃
95.	o, o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	benzyl	CH_3
96.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	3-Me-benzyl	CH ₃
97.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NHCH ₃	benzyl	H
98.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	benzyl	H
99.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	H	H
100.	o,o'-di-CH ₃ -p-CN-phenyl	F	H	CH ₃
101.	o,o'-di-CH3-p-CN-phenyl	F	benzyl	CH ₃
102.	o,o'-di-CH ₃ -p-CN-phenyl	F	benzyl	H
103.	o,o'-di-CH ₃ -p-CN-phenyl	F	H	H
104.	o,o'-diCH₃O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	Н	F
105.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	benzyl	F
106.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	benzyl	F
107.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	H	F
108.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	Н	F
109.		SO ₂ NH ₂	benzyl	F
	4-cyclopropylnaphth-l-yl	F	H	F
	4-cyclopropylnaphth-l-yl	F	benzyl	F
	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	H	F
113.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	benzyl	F
114.	o,o'-di-CH ₃ O-p-CN-phenyl	F	H	F
115.	o, o'-di-CH ₃ O-p-CN-phenyl	F	benzyl	F
116.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	H	F
117.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	benzyl	F
118.	o,o'-di-CH ₃ -p-CN-phenyl	F F	H	F F
119.	o,o'-di-CH ₃ -p-CN-phenyl		benzyl	
120.	2,4,6-trimethyl phenyl	CN	H 11	CH ₃
121.	2,4,6-trimethyl phenyl	CN	benzyl	CH ₃
122.	2,4,6-trimethyl phenyl	CN	benzyl	<u>H</u>
	2,4,6-trimethyl phenyl	CN	H	H
124.	2,4,6-trimethyl phenyl	CH=CHCN	Н	CH ₃

V Ar W Z 2,4,6-trimethyl phenyl CH=CHCN 125. CH_3 benzyl **126.** 2,4,6-trimethyl phenyl CH=CHCN benzyl Η **127.** 2,4,6-trimethyl phenyl CH=CHCN H Η 128. 2,4,6-trimethyl phenyl Η F CN 129. 2,4,6-trimethyl phenyl CN benzyl F **130.** 2,4,6-trimethyl phenyl CH=CHCN F Η 131. 2,4,6-trimethyl phenyl CH=CHCN benzyl F 132. 2,4,6-trimethyl phenyl SO₂NH₂ Η CH₃ **133.** 2,4,6-trimethyl phenyl SO₂NH₂ benzyl CH_3 2,4,6-trimethyl phenyl 134. SO₂NH₂ Η benzyl 135. 2,4,6-trimethyl phenyl SO₂NH₂ Η H \overline{H} 136. 2,4,6-trimethyl phenyl CH_3 F 137. 2,4,6-trimethyl phenyl F benzyl CH₂ 138. 2,4,6-trimethyl phenyl F benzyl Η 139. 4-cyclopropyl phenyl F \mathbf{H} Η 4-cyclopropyl phenyl SO2NH2 140. \mathbf{H} F 141. 4-cyclopropyl phenyl SO₂NH₂ benzyl F 4-cyclopropyl phenyl F F 142. Η F 143. 4-cyclopropyl phenyl benzyl F 144. o, o'-dimethyl-p-cyclopropyl phenyl CN H CH₃ 145. o, o'-dimethyl-p-cyclopropyl phenyl CN benzyl CH₃ 146. o, o'-dimethyl-p-cyclopropyl phenyl CN benzyl H 147. o, o'-dimethyl-p-cyclopropyl phenyl CN H H 148. o, o'-dimethyl-p-cyclopropyl phenyl CH=CHCN H CH₃ 149. o, o'-dimethyl-p-cyclopropyl phenyl CH=CHCN benzyl CH₃ 150. o, o'-dimethyl-p-cyclopropyl phenyl CH=CHCN benzyl H 151. o, o'-dimethyl-p-cyclopropyl phenyl CH=CHCN H Н 152. o,o'-dimethyl-p-cyclopropyl phenyl Η CN F 153. o,o'-dimethyl-p-cyclopropyl phenyl CN benzyl F CH=CHCN 154. o, o'-dimethyl-p-cyclopropyl phenyl Η F 155. o, o'-dimethyl-p-cyclopropyl phenyl CH=CHCN F benzyl 156. o,o'-dimethyl-p-cyclopropyl phenyl SO₂NH₂ Η CH_3 157. o,o'-dimethyl-p-cyclopropyl phenyl SO₂NH₂ benzyl CH₃ o,o'-dimethyl-p-cyclopropyl phenyl 158. SO₂NH₂ benzyl Η 159. o,o'-dimethyl-p-cyclopropyl phenyl SO₂NH₂ Η H 160. o,o'-dimethyl-p-cyclopropyl phenyl \mathbf{F} Η CH₃ 161. o,o'-dimethyl-p-cyclopropyl phenyl F benzyl CH₃ o, o'-dimethyl-p-cyclopropyl phenyl 162. F benzyl H o,o'-dimethyl-p-cyclopropyl phenyl F 163. H H 164. o, o'-dimethyl-p-cyclopropyl phenyl SO₂NH₂ H F SO₂NH₂ 165. o,o'-dimethyl-p-cyclopropyl phenyl benzyl F 166. o, o'-dimethyl-p-cyclopropyl phenyl F H F

10/6/2008, EAST Version: 2.3.0.3

F

benzyl

F

o,o'-dimethyl-p-cyclopropyl phenyl

167.

	Ar	V	W	Z
172.	o, o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	CH ₃	CH ₃
173.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	cyclopropyl	CH ₃
174.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	cyclopropyl	H
175.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	CH ₃	H
176.	4-cyclopropylnaphth-l-yl	CN	CH ₃	CH ₃
177.	4-cyclopropylnaphth-l-yl	CN	cyclopropyl	CH ₃
178.	4-cyclopropylnaphth-l-yl	CN	cyclopropyl	H
179.	4-cyclopropylnaphth-l-yl	CN	CH ₃	H
180.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	CH ₃
181.	4-cyclopropylnaphth-l-yl	CH=CHCN	cyclopropyl	CH ₃
182.	4-cyclopropylnaphth-l-yl	CH=CHCN	cyclopropyl	H
183.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	H
184.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	CH ₃
185.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	cyclopropyl	CH ₃
186.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	cyclopropyl	H
187.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	H
188.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	CH ₃
189.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	cyclopropyl	CH ₃
190.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	cyclopropyl	H
191.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	H
192.	o,o'-di-CH3-p-CN-phenyl	CN	CH ₃	CH_3
193.	o,o'-di-CH ₃ -p-CN-phenyl	CN	cyclopropyl	CH ₃
194.	o,o'-di-CH ₃ -p-CN-phenyl	CN	cyclopropyl	H
195.	o, o'-di-CH ₃ -p-CN-phenyl	CN	CH ₃	H
196.	o,o'-di-CH ₃ -p-CN-phenyl	CH≃CHCN	CH ₃	CH ₃
197.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	cyclopropyl	CH ₃
198.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	cyclopropyl	H
199.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃	Н
200.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	CH ₃	F
201.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	cyclopropyl	F
202.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	cyclopropyl	F
203.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	CH ₃	F
204.	4-cyclopropylnaphth-l-yl	CN	CH ₃	F
205.	4-cyclopropylnaphth-l-yl	CN	cyclopropyl	<u>F</u>
	4-cyclopropylnaphth-l-yl	CH=CHCN	CH_3	<u>F</u>
207.	4-cyclopropylnaphth-l-yl	CH=CHCN	cyclopropyl	F
208.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	F
209.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	cyclopropyl	F
210.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	F
211.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	cyclopropyl	F
212.	o, o'-di-CH ₃ -p-CN-phenyl	CN	CH ₃	F
213.	o, o'-di-CH ₃ -p-CN-phenyl	CN	cyclopropyl	F
214.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃	<u>F</u>
215.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	cyclopropyl	F
216.	o, o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	CH ₃	CH ₃
217.	o, o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	cyclopropyl	CH ₃
218.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	cyclopropyl	H

	Ar	V	W	Z
219.	o, o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	CH ₃	H
220.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NHCH ₃	CH ₃	CH ₃
221.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	cyclopropyl	CH ₃
222.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	cyclopropyl	H
223.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	CH ₃	H
224.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH ₃	CH ₃
225.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	cyclopropyl	CH ₃
226.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	cyclopropyl	H
227.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH ₃	H
228.	4-cyclopropylnaphth-l-yl	F	CH ₃	CH ₃
229.	4-cyclopropylnaphth-l-yl	F	cyclopropyl	CH ₃
230.	4-cyclopropylnaphth-l-yl	F	cyclopropyl	<u> </u>
231.	4-cyclopropylnaphth-l-yl	F	CH ₃	H
232.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	CH ₃
233.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NHCH ₃	CH ₃	CH ₃
234.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	CH ₃
235.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	<u>H</u>
236.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	H
237.	o,o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	CH ₃
238.	o,o'-di-CH ₃ O-p-CN-phenyl	F	cyclopropyl	CH ₃
239.	o,o'-di-CH ₃ O-p-CN-phenyl	F	cyclopropyl	H
240.	o,o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	Н
241.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	CH ₃	CH ₃
242.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	CH ₃
243.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	H
244.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	CH ₃	H
245.	o,o'-di-CH ₃ -p-CN-phenyl	<u>F</u>	CH ₃	CH ₃
246.	o,o'-di-CH ₃ -p-CN-phenyl	<u> </u>	cyclopropyl	CH ₃
247.	o,o'-di-CH ₃ -p-CN-phenyl	F	cyclopropyl	<u>H</u>
248.	o,o'-di-CH ₃ -p-CN-phenyl	F	CH ₃	<u>H</u>
249.	o, o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	CH ₃	<u>F</u>
250.	o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	cyclopropyl	F
251.	o, o'-diCH ₃ O-p-(CH=CHCN)phenyl	F F	cyclopropyl	<u>F</u>
252.	o, o'-diCH ₃ O-p-(CH=CHCN)phenyl	L	CH ₃	F
	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH ₃	F F
254.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂ F	cyclopropyl	
255. 256.	4-cyclopropylnaphth-l-yl 4-cyclopropylnaphth-l-yl	F	CH ₃	F F
257.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	F F
257.	o,o'-di-CN ₃ O-p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	F F
259.	o,o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	F
260.	o,o'-di-CH ₃ O-p-CN-phenyl	F	cyclopropyl	F
261.	o,o'-di-CN ₃ -p-CN-phenyl	SO ₂ NH ₂	CH ₃	F
262.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	F F
263.	o,o'-di-CH ₃ -p-CN-phenyl	F	CH ₃	F
264.	o,o'-di-CH ₃ -p-CN-phenyl	F	cyclopropyl	<u>r</u> F
265.	4-cyclopropyl phenyl	CN	CH ₃	CH ₃
200.	-r cycloptopyr phenyr	CIN	Cr13	<u> </u>

	Ar	V	W	Z
266.	2,4,6-trimethyl phenyl	CN	cyclopropyl	CH ₃
267.	2,4,6-trimethyl phenyl	CN	cyclopropyl	Н
268.	2,4,6-trimethyl phenyl	CN	CH ₃	H
269.	2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	CH ₃
270.		CH=CHCN	cyclopropyl	CH ₃
271.	2,4,6-trimethyl phenyl	CH=CHCN	cyclopropyl	Н
272.	2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	Н
273.		CN	CH ₃	F
274.		CN	cyclopropyl	F
275.		CH=CHCN	CH ₃	F
276.	2,4,6-trimethyl phenyl	CH=CHCN	cyclopropyl	F
277.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	CH ₃
278.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	cyclopropyl	CH ₃
279.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	cyclopropyl	H
280.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	Н
281.	2,4,6-trimethyl phenyl	F	CH ₃	CH ₃
282.	2,4,6-trimethyl phenyl	F	cyclopropyl	CH ₃
283.	2,4,6-trimethyl phenyl	F	cyclopropyl	H
284.	4-cyclopropyl phenyl	F	CH ₃	H
285.	4-cyclopropyl phenyl	SO ₂ NH ₂	CH ₃	F
286.	4-cyclopropyl phenyl	SO ₂ NH ₂	cyclopropyl	F
287.	4-cyclopropyl phenyl	F F	CH ₃	F
288.	4-cyclopropyl phenyl	F	cyclopropyl	F
289.	2,4,6-trimethyl phenyl	CN	CH ₃	CH ₃
290.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	cyclopropyl	CH ₃
291.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	cyclopropyl	H
292.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	CH ₃	H
293.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	·CH ₃	CH ₃
294.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	cyclopropyl	CH ₃
295.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	cyclopropyl	H
296.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	Н
297.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	CH ₃	F
298.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	cyclopropyl	F
299.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	F.
300.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	cyclopropyl	F
301.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	CH ₃	CH₃
302.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	cyclopropyl	CH ₃
303.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	cyclopropyl	H
304.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	CH ₃	H
305.	o,o'-dimethyl-p-cyclopropyl phenyl	F	CH ₃	CH ₃
306.	o,o'-dimethyl-p-cyclopropyl phenyl	F	cyclopropyl	CH ₃
307.	o,o'-dimethyl-p-cyclopropyl phenyl	F	cyclopropyl	H
308.	2,4,6-trimethyl phenyl	F	CH ₃	H
309.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	F
310.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	cyclopropyl	F
311.	2,4,6-trimethyl phenyl	F	CH ₃	F
312.	2,4,6-trimethyl phenyl	F	cyclopropyl	F

· 	Ar	, V	W	Z
313.	o,o'-di-CH3-p-acetyl-phenyl	CN	CH ₃	Н
314.	o, o'-di-CH3-p-acetyl-phenyl	CN	H	Н
315.	o,o'-di-CH3-p-acetyl-phenyl	CN	CH ₃	Cl
316.	o, o'-di-CH3-p-acetyl-phenyl	CN	H	Cl

Table 2 Contemplated Compounds of Formula IA-2

$$\bigvee_{N} \bigvee_{N} \bigvee_{N$$

	Ar	V	W	Z
1.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	F	CH ₃
2.	o o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	benzyl	CH_3
3.	o,o'diCH ₃ O-p-(CH=CHCN)-phenyl	CN	benzyl	H
4.	o,o'diCH ₃ O-p-(CH=CHCN)-phenyl	CN	F	H
5.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	Cl	CH_3
6.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	benzyl	CH ₃
7.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	benzyl	H
8.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	C1	<u>H</u>
9.	4-cyclopropylnaphth-l-yl	C≡CCH ₃	allyl	ethyl
10.	4-cyclopropylnaphth-l-yl	CN	allyl	ethyl
11.	4-cyclopropylnaphth-l-yl	CN	benzyl	H
12.	4-cyclopropylnaphth-l-yl	CN	benzyl	H
13.	4-cyclopropylnaphth-l-yl	C≡CCH ₃	allyl	ethyl
14.	4-cyclopropylnaphth-l-yl	CH=CHCN	allyl	ethyl
			3-MeO-	
15.	4-cyclopropylnaphth-l-yl	CH=CHCN	benzyl	H
16.	4-cyclopropylnaphth-l-yl	CH=CHCN	benzyl	H
17.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NHCH ₃	CH=CHCN	$\mathrm{CH_{3}}$
18.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH=CHCN	CH₃
19.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	3-Me-benzyl	H
20.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	benzyl	H
21.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH=CHCN	CH ₃
22.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₂ CH ₂ CN	CH ₃
23.	o,o'-di-CH₃O-p-CN-phenyl	CH=CHCN	CH ₂ CH ₂ CN	H
24.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	benzyl	H
25.	o,o'-di-CH ₃ O-p-(CH=CHCN)-phenyl	CN	CH ₃	H
26.	o,o'-di-CH ₃ O-p-(CH=CHCN)-phenyl	CN	CH ₃	benzyl
27.	o,o'-di-CH ₃ O-p-(CH=CHCN)-phenyl	CN	H	benzyl
28.	o,o'-di-CH ₃ O-p-(CH=CHCN)-phenyl	CN	H	H
29.	o,o'-di-CH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	CH ₃	H
30.	o,o'-di-CH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	CH ₃	benzyl
31.	o,o'-di-CH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	H	benzyl
32.	o,o'-di-CH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	Н	H
33.	4-cyclopropylnaphth-l-yl	CN	CH ₃	H
34.	4-cyclopropylnaphth-l-yl	CN	CH ₃	benzyl
35.	4-cyclopropylnaphth-l-yl	CN	Н	benzyl
36.	4-cyclopropylnaphth-l-yl	CN	Н	H

1	Ar	V	W	Z
37.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	Н
38.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	benzyl
39.	4-cyclopropylnaphth-l-yl	CH=CHCN	Н	benzyl
40.	4-cyclopropylnaphth-l-yl	CH=CHCN	H	H
41.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	Н
42.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	benzyl
43.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	Н	benzyl
44.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	Н	Н
45.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	Н
46.	o, o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	benzyl
47.	o, o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	Н	benzyl
48.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	Н	H
49.	o,o'-di-CH ₃ -p-CN-phenyl	CN	CH ₃	Н
50.	o,o'-di-CH ₃ -p-CN-phenyl	CN	CH ₃	benzyl
51.	o,o'-di-CH ₃ -p-CN-phenyl	CN	Н	benzyl
52.	o,o'-di-CH ₃ -p-CN-phenyl	CN	Н	Н
53.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃	Н
54.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃	benzyl
55.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	Н	benzyl
56.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	Н	H
57.	o, o'-di-CH ₃ O-p-(CH=CHCN)-phenyl	CN	F	Н
58.	o,o'-di-CH ₃ O-p-(CH=CHCN)-phenyl	CN	F	benzyl
59.	o,o'-di-CH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	F	benzyl
60.	o,o'-di-CH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	F	H
61.	4-cyclopropylnaphth-l-yl	CN	F	Н
62.	4-cyclopropylnaphth-l-yl	CN	F	benzyl
63.	4-cyclopropylnaphth-l-yl	CH=CHCN	F	H
64.	4-cyclopropylnaphth-l-yl	CH=CHCN	F	benzyl
65.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	F	H
66.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	F	benzyl
67.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	F	H
68.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	F	benzyl
69.	o,o'-di-CH ₃ -p-CN-phenyl	CN	F	H
70.	o,o'-di-CH ₃ -p-CN-phenyl	CN	F	benzyl
71.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	<u> </u>	H
72.	o, o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	F	benzyl
73.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	H
74.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	benzyl
75.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	H	benzyl
76.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	H	H
77.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	H
78.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	CH ₃	benzyl
79.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	H	benzyl
80.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	H	H
81.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH ₃	H
82.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH ₃	benzyl
83.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	<u>H</u>	benzyl

	Ar	V	W	Z
84.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	Н	H
85.	4-cyclopropylnaphth-l-yl	F	CH ₃	H
86.	4-cyclopropylnaphth-l-yl	F	CH ₃	benzyl
87.	4-cyclopropylnaphth-l-yl	F	H	benzyl
88.	4-cyclopropylnaphth-l-yl	F	H	Н
89.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	H
90.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	benzyl
91.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	Н	benzyl
92.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	H	Н
93.	o,o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	Н
94.	o,o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	benzyl
95.	o,o'-di-CH ₃ O-p-CN-phenyl	F	H	benzyl
96.	o,o'-di-CH ₃ O-p-CN-phenyl	F	Н	H
97.	o,o'-di-CH₃-p-CN-phenyl	SO ₂ NH ₂	CH ₃	H
98.	<i>o,o'-di-</i> CH₃- <i>p</i> -CN-phenyl	SO ₂ NH ₂	CH ₃	benzyl
99.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	H	benzyl
100.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	Н	Н
101.	o,o'-di-CH ₃ -p-CN-phenyl	F	CH ₃	H
102.	o,o'-di-CH ₃ -p-CN-phenyl	F	CH ₃	benzyl
. 103.	o,o'-di-CH ₃ -p-CN-phenyl	F	.Н	benzyl
104.	o,o'-di-CH ₃ -p-CN-phenyl	F	H	H
105.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	F	Н
106.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	F	benzyl
107.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	F	benzyl
108.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	·F	F	H
109.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	F	Н
110.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	F	benzyl
111.	4-cyclopropylnaphth-l-yl	<u> </u>	F	H
112.	4-cyclopropylnaphth-1-yl	F	F	benzyl
113.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	F	H
114.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	F	benzyl
115.	o,o'-di-CH ₃ O-p-CN-phenyl	F	F	H
116.	o,o'-di-CH ₃ O-p-CN-phenyl	F	F	benzyl
117.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	F	H
118.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	F	benzyl
119.	o,o'-di-CH ₃ -p-CN-phenyl	<u> </u>	F	H
120.	o,o'-di-CH ₃ -p-CN-phenyl	F	F	benzyl
121.	4-cyclopropyl phenyl	CN	CH ₃	H
122.	4-cyclopropyl phenyl	CN	CH ₃	benzyl
123.	4-cyclopropyl phenyl	CN	H	benzyl
124.	4-cyclopropyl phenyl	CN	H	H
125.	2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	H
126.	2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	benzyl
127.	2,4,6-trimethyl phenyl	CH=CHCN	H	benzyl
128.	2,4,6-trimethyl phenyl	CH=CHCN	H	Н
129.	2,4,6-trimethyl phenyl	CN	F	H
130.	2,4,6-trimethyl phenyl	CN	F	benzyl

	Ar	V	W	Z
131.	2,4,6-trimethyl phenyl	CH=CHCN	F	H
132.	2,4,6-trimethyl phenyl	CH=CHCN	F	benzyl
133.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	H
134.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	benzyl
135.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	Н	benzyl
136.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	H	Н
137.	2,4,6-trimethyl phenyl	F	CH ₃	H
138.	2,4,6-trimethyl phenyl	F	CH ₃	benzyl
139.	2,4,6-trimethyl phenyl	F	H	benzyl
140.	4-cyclopropyl phenyl	F	H	H
141.	4-cyclopropyl phenyl	SO ₂ NH ₂	F	H
142.	4-cyclopropyl phenyl	SO ₂ NH ₂	F	benzyl
143.	4-cyclopropyl phenyl	F	F	Н
144.	4-cyclopropyl phenyl	F	F	benzyl
145.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	CH ₃	H
146.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	CH ₃	benzyl
147.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	H	benzyl
148.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	H	H
149.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	Н
150.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	benzyl
151.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	H	benzyl
152.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	H	H
153.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	F	Н
154.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	F	benzyl
155.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	F	H
156.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	F	benzyl
157.	o,o'-dimethyl-p-cyclopropyl phenyl	SO_2NH_2	CH ₃	H
158.	o,o'-dimethyl-p-cyclopropyl phenyl	SO_2NH_2	$\mathrm{CH_{3}}$	benzyl
159.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	Н	benzyl
160.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	Н	Н
161.	o,o'-dimethyl-p-cyclopropyl phenyl	F	CH ₃	H
162.	o,o'-dimethyl-p-cyclopropyl phenyl	F	CH_3	benzyl
163.	o,o'-dimethyl-p-cyclopropyl phenyl	F	H	benzyl
164.	o,o'-dimethyl-p-cyclopropyl phenyl	F	H	H
165.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	F	H
166.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	F	benzyl
167.	2-methyl-4-cyclopropyl phenyl	F	F	H
168.	2-methyl-4-cyclopropyl phenyl	F	F	benzyl
169.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	CH ₃	CH ₃
170.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	CH ₃	cyclopropyl
171.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	H	cyclopropyl
172.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	H	CH ₃
173.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	CH ₃	CH ₃
174.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	CH ₃	cyclopropyl
175.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	H	cyclopropyl
176.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	H	CH ₃
177.	4-cyclopropylnaphth-l-yl	CN	CH ₃	CH ₃

	Ar	V	W	Z
178.	4-cyclopropylnaphth-l-yl	CN	CH ₃	cyclopropyl
179.	4-cyclopropylnaphth-l-yl	CN	H	cyclopropyl
180.	4-cyclopropylnaphth-l-yl	CN	H	CH ₃
181.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	CH ₃
182.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	cyclopropyl
183.	4-cyclopropylnaphth-1-yl	CH=CHCN	H	cyclopropyl
184.	4-cyclopropylnaphth-l-yl	CH=CHCN	H	CH ₃
185.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	CH ₃
186.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	cyclopropyl
187.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	H	cyclopropyl
188.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	H	CH ₃
189.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	CH ₃
190.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	cyclopropyl
191.	o, o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	Н	cyclopropyl
192.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	H	CH ₃
193.	o,o'-di-CH ₃ -p-CN-phenyl	CN	CH ₃	CH ₃
194.	o,o'-di-CH ₃ -p-CN-phenyl	CN	CH ₃	cyclopropyl
195.	o,o'-di-CH ₃ -p-CN-phenyl	CN	H	cyclopropyl
196.	o,o'-di-CH ₃ -p-CN-phenyl	CN	H	CH ₃
197.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃	CH ₃
198.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃	cyclopropyl
199.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	Н	cyclopropyl
200.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	H	CH ₃
201.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	F	CH ₃
202.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	F	cyclopropyl
203.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	F	cyclopropyl
204.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	F	CH_3
205.	4-cyclopropylnaphth-l-yl	CN	F	CH ₃
206.	4-cyclopropylnaphth-l-yl	CN	F	cyclopropyl
207.	4-cyclopropylnaphth-l-yl	CH=CHCN	F	CH ₃
208.	4-cyclopropylnaphth-l-yl	CH=CHCN	F	cyclopropyl
209.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	F	CH ₃
210.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	F	cyclopropyl
211.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	F	CH ₃
212.	o,o'-di-CH₃O-p-CN-phenyl	CH=CHCN	F	cyclopropyl
213.	o,o'-di-CH ₃ -p-CN-phenyl	CN	F	CH ₃
214.	o,o'-di-CH ₃ -p-CN-phenyl	CN	F	cyclopropyl
215.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	F	CH ₃
216.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	F	cyclopropyl
217.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	CH ₃
218.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	cyclopropyl
219.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	H	cyclopropyl
220.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	H	CH ₃
221.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	CH ₃
222.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	<u> </u>	CH ₃	cyclopropyl
223.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	H	cyclopropyl
224.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	<u> </u>	CH ₃

	Ar	V	W	Z
225.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH ₃	CH ₃
226.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH ₃	cyclopropyl
227.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	H	cyclopropyl
228.	4-cyclopropylnaphth-I-yl	SO ₂ NH ₂	H	CH ₃
229.	4-cyclopropylnaphth-l-yl	F	CH ₃	CH ₃
. 230.	4-cyclopropylnaphth-l-yl	F	CH ₃	cyclopropyl
231.	4-cyclopropylnaphth-l-yl	F	H	cyclopropyl
232.	4-cyclopropylnaphth-l-yl	F	H	CH ₃
233.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	CH ₃
234.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	cyclopropyl
235.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	H	cyclopropyl
236.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	H	CH ₃
237.	o,o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	CH ₃
238.	o,o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	cyclopropyl
239.	o,o'-di-CH ₃ O-p-CN-phenyl	F	H	cyclopropyl
240.	o,o'-di-CH ₃ O-p-CN-phenyl	F ·	H	CH ₃
241.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	CH ₃	CH ₃
242.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	CH ₃	cyclopropyl
243.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	H	cyclopropyl
244.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	H	CH ₃
245.	o,o'-di-CH ₃ -p-CN-phenyl	F	CH ₃	CH ₃
246.	o,o'-di-CH ₃ -p-CN-phenyl	F	CH ₃	cyclopropyl
247.	o,o'-di-CH ₃ -p-CN-phenyl	F	H	cyclopropyl
248.	o,o'-di-CH ₃ -p-CN-phenyl	F	H	CH ₃
249.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	F	CH ₃
250.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	F	cyclopropyl
251.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	F	cyclopropyl
252.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	F	CH ₃
253.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	F	CH ₃
254.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	F	cyclopropyl
255.	4-cyclopropylnaphth-l-yl	F	F	CH ₃
256.	4-cyclopropylnaphth-l-yl	F	F	cyclopropyl
257.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	F	CH ₃
258.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	F	cyclopropyl
259.	o,o'-di-CH ₃ O-p-CN-phenyl	F	F	CH ₃
260.	o,o'-di-CH ₃ O-p-CN-phenyl	F	F	cyclopropyl
261.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	F	CH ₃
262.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	F	cyclopropyl
263.	o,o'-di-CH ₃ -p-CN-phenyl	F	F	CH ₃
264.	o,o'-di-CH ₃ -p-CN-phenyl	F	F	cyclopropyl
265.	4-cyclopropyl phenyl	CN	CH ₃	CH ₃
266.	2,4,6-trimethyl phenyl	CN	CH ₃	cyclopropyl
267.	2,4,6-trimethyl phenyl	CN	H	cyclopropyl
268.	2,4,6-trimethyl phenyl	CN	H	CH ₃
269.	2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	CH ₃
270.	2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	cyclopropyl
271.	2,4,6-trimethyl phenyl	CH=CHCN	H	cyclopropyl

	Ar	V	· W	Z
272.	2,4,6-trimethyl phenyl	CH=CHCN	H	CH ₃
273.	2,4,6-trimethyl phenyl	CN	F	CH ₃
274.	2,4,6-trimethyl phenyl	CN	F	cyclopropyl
275.	2,4,6-trimethyl phenyl	CH=CHCN	F	CH ₃
276.	2,4,6-trimethyl phenyl	CH=CHCN	F	cyclopropyl
277.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	CH ₃
278.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	cyclopropyl
279.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	Ĥ	cyclopropyl
280.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	H	CH ₃
281.	2,4,6-trimethyl phenyl	F	CH ₃	CH ₃
282.	2,4,6-trimethyl phenyl	F	CH ₃	cyclopropyl
283.	2,4,6-trimethyl phenyl	F	Н	cyclopropyl
284.	2,4,6-trimethyl phenyl	F	H	CH ₃
285.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	F	CH ₃
286.	4-cyclopropyl phenyl	SO ₂ NH ₂	F	cyclopropyl
287.	4-cyclopropyl phenyl	F	F	CH ₃
288.	4-cyclopropyl phenyl	F	F	cyclopropyl
289.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	CH ₃	CH ₃
290.	o, o'-dimethyl-p-cyclopropyl phenyl	CN	CH ₃	cyclopropyl
291.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	H	cyclopropyl
292.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	H	CH ₃
293.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	CH ₃
294.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	cyclopropyl
295.	o, o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	H	cyclopropyl
296.	o, o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	H	CH ₃
297.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	F	CH ₃
298.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	F	cyclopropyl
299.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	F	CH ₃
300.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	F	cyclopropyl
301.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	CH ₃	CH ₃
302.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	CH ₃	cyclopropyl
303.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	Н	cyclopropyl
304.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	H	CH ₃
305.	o,o'-dimethyl-p-cyclopropyl phenyl	F	CH ₃	CH ₃
306.	o,o'-dimethyl-p-cyclopropyl phenyl	F	CH ₃	cyclopropyl
307.	o,o'-dimethyl-p-cyclopropyl phenyl	F	H	cyclopropyl
308.	o,o'-dimethyl-p-cyclopropyl phenyl	F	H	CH ₃
309.	o,o'-dimethyl-p-cyclopropyl phenyl	SO_2NH_2	F	CH ₃
310.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	F	cyclopropyl
311.	o,o'-dimethyl-p-cyclopropyl phenyl	F	F	CH ₃
312.	o,o'-dimethyl-p-cyclopropyl phenyl	F	F	cyclopropyl
313.	o,o'-di-CH ₃ -p-acetyl-phenyl	CN	H	H
314.	o,o'-di-CH ₃ -p-acetyl-phenyl	CN	CH ₃	Н
315.	o,o'-di-CH ₃ -p-acetyl-phenyl	CN	H	Cl
316.	o,o'-di-CH ₃ -p-acetyl-phenyl	CN	CH ₃	CI

Table 3 Contemplated Compounds of Formula IA-3

	Ar	V	W	Z
1.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	benzyl	F
2.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	benzyl	C1
3.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	allyl	F
4.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	allyl	C1
5.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	benzyl	CH ₃
		"s	3-MeO-	
6.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	benzyl	CH ₃
			3-Me-	
7.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	benzyl	CH ₃
8.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	allyl	CH ₃
9.	4-cyclopropylnaphth-l-yl	CN	CH₂CH₃	H
10.	4-cyclopropylnaphth-l-yl	CN	isopropyl	Н
11.	4-cyclopropylnaphth-l-yl	CN	CH ₂ CF ₃	Br
12.	4-cyclopropylnaphth-l-yl	CN	CH ₂ CF ₃	C1
13.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₂ CH ₃	H
14.	4-cyclopropylnaphth-1-yl	CH=CHCN	CH ₂ CH ₃	Br
15.	4-cyclopropylnaphth-1-yl	CH=CHCN	CH ₂ CF ₃	CH ₃
16.	4-cyclopropylnaphth-1-yl	CH=CHCN	CH ₂ CF ₃	H
17.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	benzyl	F
18.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	benzyl	Cl
19.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	allyl	F
20.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	allyl	C1
21.	o,o'-di-CH3O-p-CN-phenyl	CH=CHCN	benzyl	CH ₃
22.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	benzyl	Br
23.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	allyl	CH ₃
24.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	allyl	H
25.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	H	F
26.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	benzyl	C1
27.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	benzyl	H
28.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	H	H
29.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	H	CH ₃
30.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	benzyl	CH ₃
31.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	benzyl	H
32.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	H	H
33.	4-cyclopropylnaphth-l-yl	CN	H	CH ₃
34.	4-cyclopropylnaphth-l-yl	CN	benzyl	CH ₃
35.	4-cyclopropylnaphth-l-yl	CN	benzyl	H

	Ar	V	W	Z
36.	4-cyclopropylnaphth-l-yl	CN	Н	Н
37.	4-cyclopropylnaphth-l-yl	CH=CHCN	Н	CH ₃
38.	4-cyclopropylnaphth-l-yl	CH=CHCN	benzyl	CH ₃
39.	4-cyclopropylnaphth-l-yl	CH=CHCN	benzyl	H
40.	4-cyclopropylnaphth-l-yl	CH=CHCN	H	H
41.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	H	CH ₃
42.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	benzyl	CH ₃
43.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	benzyl	H
44.	o,o'-di-CH₃O-p-CN-phenyl	CN	H	H
45.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	H	CH ₃
46.	o,o'-di-CH₃O-p-CN-phenyl	CH=CHCN	benzyl	CH ₃
. 47.	o,o'-di-CH₃O-p-CN-phenyl	CH=CHCN	benzyl	H
48.	o,o'-di-CH₃O-p-CN-phenyl	CH=CHCN	H	Н
49.	o,o'-di-CH ₃ -p-CN-phenyl	CN	H	CH ₃
50.	o,o'-di-CH3-p-CN-phenyl	CN	benzyl	CH ₃
51.	o,o'-di-CH3-p-CN-phenyl	CN	benzyl	H
52.	o,o'-di-CH ₃ -p-CN-phenyl	CN	H	H
53.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	Н	CH ₃
54.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	benzyl	CH ₃
55.	o, o'- di -CH ₃ - p -CN-phenyl	CH=CHCN	benzyl	. H
56.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	Н	H
57.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	H	F
58.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	benzyl	F
59.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	benzyl	F
60.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	H	F
61.	4-cyclopropylnaphth-l-yl	CN	H	F
62.	4-cyclopropylnaphth-l-yl	CN	benzyl	F
63.	4-cyclopropylnaphth-l-yl	CH=CHCN	H	F
64.	4-cyclopropylnaphth-l-yl	CH=CHCN	benzyl	F
65.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	H	F
66.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	benzyl	F
67.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	H	F
68.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	benzyl	F
69.	o,o'-di-CH ₃ -p-CN-phenyl	CN	H	F
70.	o,o'-di-CH ₃ -p-CN-phenyl	CN	benzyl	F
71.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	H	F
72.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	benzyl	F
73.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	H	CH ₃
74.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	benzyl	CH ₃
75.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	benzyl	H
76.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	H	H
77.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	H	CH ₃
78.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	benzyl	CH ₃
79.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	benzyl	H
80.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	H	H
81.	4-cyclopropylnaphth-1-yl	SO ₂ NH ₂	H	CH ₃
82.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	benzyl	CH ₃

	Ar	V	W	Z
83.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	benzyl	H
84.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	H	H
85.	4-cyclopropylnaphth-l-yl	F	Н	CH ₃
86.	4-cyclopropylnaphth-l-yl	F	benzyl	CH ₃
87.	4-cyclopropylnaphth-1-yl	F	benzyl	Н
88.	4-cyclopropylnaphth-l-yl	F	H	H
89.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	Н	CH ₃
90.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	benzyl	CH ₃
91.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	benzyl	H
92.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	H	Н
93.	o,o'-di-CH ₃ O-p-CN-phenyl	F	Н	CH ₃
94.	o,o'-di-CH ₃ O-p-CN-phenyl	F	benzyl	CH ₃
95.	o,o'-di-CH ₃ O-p-CN-phenyl	F	benzyl	Н
96.	o,o'-di-CH ₃ O-p-CN-phenyl	F	H	Н
97.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	H	CH ₃
98.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	benzyl	CH ₃
99.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	benzyl	H
100.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	H	Н
101.	o,o'-di-CH ₃ -p-CN-phenyl	F	H	CH ₃
102.	o,o'-di-CH ₃ -p-CN-phenyl	F	benzyl	CH ₃
103.	o,o'-di-CH ₃ -p-CN-phenyl	F	benzyl	H
104.	o,o'-di-CH ₃ -p-CN-phenyl	F	H	H
105.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	H	F
106.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	benzyl	F
107.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	benzyl	F
108.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	H	F
109.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	H	F
110.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	benzyl	F
111.	4-cyclopropylnaphth-l-yl	F	H	F
112.	4-cyclopropylnaphth-l-yl	F	benzyl	F
113.	o,o'-di-CH ₃ O-p-CN-phenyl	SO_2NH_2	H	F
114.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	benzyl	F
115.	o,o'-di-CH ₃ O-p-CN-phenyl	F	H	F
116.	o,o'-di-CH ₃ O-p-CN-phenyl	F	benzyl	F
117.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	H	F
118.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	benzyl	F
119.	o,o'-di-CH ₃ -p-CN-phenyl	F	H	F
120.	o,o'-di-CH ₃ -p-CN-phenyl	F	benzyl	F
121.	2,4,6-trimethyl phenyl	CN	H	CH ₃
122.	2,4,6-trimethyl phenyl	CN	benzyl	CH ₃
123.	2,4,6-trimethyl phenyl	CN	benzyl	Н
124.	2,4,6-trimethyl phenyl	CN	H	H
125.	2,4,6-trimethyl phenyl	CH=CHCN	H	CH ₃
126.	2,4,6-trimethyl phenyl	CH=CHCN	benzyl	CH ₃
127.	4-cyclopropyl phenyl	CH=CHCN	benzyl	H
128.	4-cyclopropyl phenyl	CH=CHCN	H	H
129.	4-cyclopropyl phenyl	CN	H	F

	Ar	V	W	Z
130.	4-cyclopropyl phenyl	CN	benzyl	F
131.	4-cyclopropyl phenyl	CH=CHCN	H	F
132.	2,4,6-trimethyl phenyl	CH=CHCN	benzyl	F
133.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	H	CH ₃
134.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	benzyl	CH ₃
135.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	benzyl	Н
136.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	H	Н
137.	2,4,6-trimethyl phenyl	F	H	CH ₃
138.	2,4,6-trimethyl phenyl	F	benzyl	CH ₃
139.	2,4,6-trimethyl phenyl	F	benzyl	H
140.	2,4,6-trimethyl phenyl	F	H	H
141.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	Н	F
142.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	benzyl	F
143.	2,4,6-trimethyl phenyl	F	Н	F
144.	2,4,6-trimethyl phenyl	F	benzyl	F
145.	2,4,6-trimethyl phenyl	CN	H	CH ₃
146.	2,4,6-trimethyl phenyl	CN	benzyl	CH ₃
147.	2,4,6-trimethyl phenyl	CN	benzyl	H
148.	2,4,6-trimethyl phenyl	CN	H	H
149.	2,4,6-trimethyl phenyl	CH=CHCN	H	ÇH₃
150.	2,4,6-trimethyl phenyl	CH=CHCN	benzyl	CH ₃
151.	2,4,6-trimethyl phenyl	CH=CHCN	benzyl	H
152.	2,4,6-trimethyl phenyl	CH=CHCN	H	H
153.	2,4,6-trimethyl phenyl	CN	H	F
154.	2,4,6-trimethyl phenyl	CN	benzyl	F
155.	2,4,6-trimethyl phenyl	CH=CHCN	H	F
156.	2,4,6-trimethyl phenyl	CH=CHCN	benzyl	F
157.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	H	CH ₃
158.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	benzyl	$_{\rm CH_3}$
159.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	benzyl	H
160.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	H	H
161.	o,o'-dimethyl-p-cyclopropyl phenyl	F	H	CH ₃
162.	o,o'-dimethyl-p-cyclopropyl phenyl	F	benzyl	CH ₃
163.	o,o'-dimethyl-p-cyclopropyl phenyl	F	benzyl	H
164.	o,o'-dimethyl-p-cyclopropyl phenyl	F	Н	H
165.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	H	F
166.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	benzyl	F
167.	o, o'-dimethyl-p-cyclopropyl phenyl	F	H	F
168. 169.	o, o'-dimethyl-p-cyclopropyl phenyl	F	benzyl	F
	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	CH ₃	CH ₃
170. 171.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	cyclopropyl	CH ₃
171.	o, o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	cyclopropyl	H
172.	o, o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH-CHCM	CH ₃	H
	o, o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	CH ₃	CH ₃
174. 175.	o, o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	cyclopropyl	CH ₃
176.	o, o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	cyclopropyl	<u>H</u>
1/0.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	CH ₃	H

	Ar	V	W	Z
177.	4-cyclopropylnaphth-l-yl	CN	CH ₃	CH ₃
178.	4-cyclopropylnaphth-l-yl	CN	cyclopropyl	CH ₃
179.	4-cyclopropylnaphth-l-yl	CN	cyclopropyl	H
180.	4-cyclopropylnaphth-l-yl	CN	CH ₃	Н
181.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	CH ₃
182.	4-cyclopropylnaphth-l-yl	CH=CHCN	cyclopropyl	CH ₃
183.	4-cyclopropylnaphth-l-yl	CH=CHCN	cyclopropyl	H
184.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	H
185.	o,o'-di-CH₃O-p-CN-phenyl	CN	CH ₃	CH ₃
186.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	cyclopropyl	CH ₃
187.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	cyclopropyl	H
188.	<i>o,o′-di</i> -CH₃O- <i>p</i> -CN-phenyl	CN	CH ₃	H
189.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	CH ₃
190.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	cyclopropyl	CH ₃
191.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	cyclopropyl	Н
192.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	H
193.	o,o'-di-CH ₃ -p-CN-phenyl	CN	CH ₃	CH ₃
194.	o,o'-di-CH ₃ -p-CN-phenyl	CN	cyclopropyl	. CH ₃
195.	o,o'-di-CH ₃ -p-CN-phenyl	CN	cyclopropyl	Н
196.	o,o'-di-CH ₃ -p-CN-phenyl .	CN	CH ₃	Н
197.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃	CH ₃
198.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	cyclopropyl	CH ₃
199.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	cyclopropyl	H
200.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃	H
201.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	CH ₃	F
202.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	cyclopropyl	F
203.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	cyclopropyl	F
204. 205.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	CH ₃	F
	4-cyclopropylnaphth-l-yl	CN	CH ₃	F
206. 207.	4-cyclopropylnaphth-1-yl	CN	cyclopropyl	F
208.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	F
208.	4-cyclopropylnaphth-l-yl	CH=CHCN	cyclopropyl	F
210.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	F
210.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	cyclopropyl	F
212.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	<u>F</u>
213.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	cyclopropyl	F
214.	o,o'-di-CH ₃ -p-CN-phenyl	CN	CH ₃	F
215.		CH-CHCN	cyclopropyl	F
216.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃	F
217.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	cyclopropyl	F GY
218.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	CH ₃
219.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	cyclopropyl	CH ₃
220.	o,o'-diCH ₂ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	cyclopropyl	H
221.	o,o'-diCH ₂ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	H
222.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	CH ₃
223.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	cyclopropyl	CH ₃
443.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	cyclopropyl	H

	Ar	V	W	Z
224.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	CH ₃	Н
225.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH ₃	CH ₃
226.	4-cyclopropylnaphth-1-yl	SO ₂ NH ₂	cyclopropyl	CH ₃
227.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	cyclopropyl	H
228.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH ₃	H
229.	4-cyclopropylnaphth-l-yl	F	CH ₃	CH ₃
230.	4-cyclopropylnaphth-l-yl	F	cyclopropyl	CH ₃
231.	4-cyclopropylnaphth-1-yl	F	cyclopropyl	H
232.	4-cyclopropylnaphth-l-yl	F	CH ₃	H
233.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	CH ₃
234.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	CH ₃
235.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	. H
236.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	H
237.	o,o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	CH ₃
238.	o,o'-di-CH ₃ O-p-CN-phenyl	F	cyclopropyl	CH ₃
239.	o,o'-di-CH ₃ O-p-CN-phenyl	F	cyclopropyl	H
240.	o,o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	H
241.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	CH ₃	CH ₃
242.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	CH ₃
243.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	H
244.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	CH ₃	H
245.	o,o'-di-CH ₃ -p-CN-phenyl	F	CH ₃	CH ₃
246.	o,o'-di-CH ₃ -p-CN-phenyl	F	cyclopropyl	CH ₃
247.	o,o'-di-CH ₃ -p-CN-phenyl	F	cyclopropyl	H
248.	o,o'-di-CH ₃ -p-CN-phenyl	F	CH ₃	H
249.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	F
250.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	cyclopropyl	F
251.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	cyclopropyl	F
252.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	CH ₃	F
253.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH ₃	F
254.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	cyclopropyl	F
255.	4-cyclopropylnaphth-l-yl	F	CH ₃	F
256.	4-cyclopropylnaphth-l-yl	F	cyclopropyl	F
257.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	F
258.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	F
259.	o,o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	F
260.	o,o'-di-CH ₃ O-p-CN-phenyl	F	cyclopropyl	F
261.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	CH ₃	F
262.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	F
263.	o,o'-di-CH3-p-CN-phenyl	F	CH ₃	F
264.	o,o'-di-CH ₃ -p-CN-phenyl	F	cyclopropyl	F
265.	4-cyclopropyl phenyl	CN	CH ₃	CH ₃
266.	4-cyclopropyl phenyl	CN	cyclopropyl	CH ₃
267.	2,4,6-trimethyl phenyl	CN	cyclopropyl	H
268.	2,4,6-trimethyl phenyl	CN	CH ₃	H
269.	2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	CH ₃
270.	2,4,6-trimethyl phenyl	CH=CHCN	cyclopropyl	$-\frac{\text{CH}_3}{\text{CH}_3}$

	Ar	V	W	Z
271.	2,4,6-trimethyl phenyl	CH=CHCN	cyclopropyl	H
272.	2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	H
273.	2,4,6-trimethyl phenyl	CN	CH ₃	F
274.	2,4,6-trimethyl phenyl	CN	cyclopropyl	F
275.	2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	F
276.	2,4,6-trimethyl phenyl	CH=CHCN	cyclopropyl	F
277.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	CH ₃
278.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	cyclopropyl	CH ₃
279.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	cyclopropyl	H
280.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	H
281.	2,4,6-trimethyl phenyl	F	CH ₃	CH_3
282.	2,4,6-trimethyl phenyl	F	cyclopropyl	CH_3
283.	2,4,6-trimethyl phenyl	F	cyclopropyl	H
284.	2,4,6-trimethyl phenyl	F	CH ₃	Н
285.	4-cyclopropyl phenyl	SO ₂ NH ₂	CH ₃	F
286.	4-cyclopropyl phenyl	SO ₂ NH ₂	cyclopropyl	F
287.	4-cyclopropyl phenyl	F	CH ₃	F
288.	4-cyclopropyl phenyl	F	cyclopropyl	F
289.	2,4,6-trimethyl phenyl	CN	CH ₃	CH ₃
290.	2,4,6-trimethyl phenyl	CN	cyclopropyl	CH ₃
291.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	cyclopropyl	H
292.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	CH ₃	Н
293.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	CH ₃
294.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	cyclopropyl	CH ₃
295.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	cyclopropyl	<u>H</u>
296.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	<u>H</u>
297.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	CH ₃	F
298.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	cyclopropyl	F
299.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	F
300.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	cyclopropyl	F
301.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	CH ₃	CH ₃
302.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	cyclopropyl	CH ₃
303.	o, o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	cyclopropyl	H
304.	o,o'-dimethyl-p-cyclopropyl phenyl o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	CH ₃ CH ₃	H CH ₃
305.	o, o'-dimethyl-p-cyclopropyl phenyl	F		
307.	o, o'-dimethyl-p-cyclopropyl phenyl	F	cyclopropyl	CH ₃
308.	o, o'-dimethyl-p-cyclopropyl phenyl	F	cyclopropyl	H H
309.	2,4,6-trimethyl phenyl		CH ₃ CH ₃	
310.	2,4,6-trimethyl phenyl	SO ₂ NH ₂ SO ₂ NH ₂		F F
310.	2,4,6-trimethyl phenyl	F SO ₂ NH ₂	cyclopropyl CH ₃	F F
311.	2,4,6-trimethyl phenyl	F		F
312.	o,o'-di-CH ₃ -p-acetyl-phenyl	CN	cyclopropyl H	H
314.	o,o'-di-CH ₃ -p-acetyl-phenyl	CN	CH ₃	H
314.	o, o'-di-CH ₃ -p-acetyl-phenyl	CN	H	Cl
316.	o,o'-di-CH ₃ -p-acetyl-phenyl	CN	CH ₃	Cl
2101	$0,0$ - α -C113- ρ -accety1-pheny1	CIA	СПЗ	CI

Table 4 Contemplated Compounds of Formula IA-4

5

<u> </u>	Ar	V	W	Z
1.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	Н	H
2.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	CH ₃	H
3.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	F	CH ₃
4.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	C1	CH ₃
5.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	F	H
6.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	Cl	H
7.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	Br	CH ₃
8.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	H	CH ₃
9.	4-cyclopropylnaphth-l-yl	CN	H	Н
10.	4-cyclopropylnaphth-l-yl	CN	CH ₃	H
11.	4-cyclopropylnaphth-l-yl	CN	F	CH ₃
12.	4-cyclopropylnaphth-l-yl	CN	C1	CH ₃
13.	4-cyclopropylnaphth-l-yl	CH=CHCN	C1	H
14.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	Н
15.	4-cyclopropylnaphth-l-yl	CH=CHCN	F	CH ₃
16.	4-cyclopropylnaphth-l-yl	CH=CHCN	C1	CH ₃
17.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	Br	H
18.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	H	Н
19.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	CH ₃
20.	o,o'-di-CH₃O-p-CN-phenyl	CN	F	CH ₃
21.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	Н	H
22.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	Н
23.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	F	CH ₃
24.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	C1	CH ₃
25.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	F	CH ₃
26.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	C1	CH ₃
27.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	C1	Н
28.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	F	H
29.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	H	CH ₃
30.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	CH ₃	CH ₃
31.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	F	H
32.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	C1	H
33.	4-cyclopropylnaphth-l-yl	CN	H	ethyl
34.	4-cyclopropylnaphth-l-yl	CN	C1	ethyl
35.	4-cyclopropylnaphth-l-yl	CN	F	H
36.	4-cyclopropylnaphth-l-yl	CN	Cl	H

	Ar	V	W	Z
37.	4-cyclopropylnaphth-l-yl	CH=CHCN	Br	ethyl
38.	4-cyclopropylnaphth-l-yl	CH=CHCN	H	ethyl
39.	4-cyclopropylnaphth-l-yl	CH=CHCN	H	H
40.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	H
41.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	F	CH ₃
42.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	Cl	. CH ₃
43.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	Cl	H H
44.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	F	H
45.	o,o'-di-CH ₃ O-p-CN-phenyl	C≡CCH ₃	Cl	CH ₃
46.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	Br	CH ₃
47.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	Br	H
48.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	H	H
49.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	CH ₃	H
50.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	CH ₃	benzyl
51.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	H	benzyl
52.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	, H	H
53.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	CH ₃	H
54.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	CH ₃	benzyl
55.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	H	benzyl
56.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	H	H
57.	4-cyclopropylnaphth-l-yl	CN	CH ₃	H
58.	4-cyclopropylnaphth-l-yl	CN	CH ₃	benzyl
59.	4-cyclopropylnaphth-l-yl	CN	H	benzyl
60.	4-cyclopropylnaphth-l-yl	CN	H	H
61.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	H
62.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	benzyl
63.	4-cyclopropylnaphth-l-yl	CH=CHCN	H	benzyl
54.	4-cyclopropylnaphth-l-yl	CH=CHCN	H	H
65.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	H
66.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	benzyl
57.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	H	benzyl
58.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	Н	H
59.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	H
70.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	benzyl
71.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	Н	benzyl
72.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	H	H
73.	o,o'-di-CH ₃ -p-CN-phenyl	CN	CH ₃	H
74.	o,o'-di-CH ₃ -p-CN-phenyl	CN	CH ₃	benzyl
75.	o,o'-di-CH ₃ -p-CN-phenyl	CN	Н	benzyl
76.	o,o'-di-CH ₃ -p-CN-phenyl	CN	H	H
77.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃	H
8.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃	benzyl
9.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	H	benzyl
0.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	H	H
31.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	F	H
2.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	F	benzyl
3.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	F	benzyl

	Ar	V	W	Z
84.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	F	Н
85.	4-cyclopropylnaphth-l-yl	CN	F	Н
86.	4-cyclopropylnaphth-l-yl	CN	F	benzyl
87.	4-cyclopropylnaphth-l-yl	CH=CHCN	F	H
88.	4-cyclopropylnaphth-l-yl	CH=CHCN	F	benzyl
89.	o,o'-di-CH ₃ O-p-CN-phenyl	. CN	F	Н
90.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	F	benzyl
91.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	F	H
92.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	F	benzyl
93.	o,o'-di-CH ₃ -p-CN-phenyl	CN	·F	Н
94.	o,o'-di-CH ₃ -p-CN-phenyl	CN	F	benzyl
95.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	F	H
96.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	F	benzyl
97.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	H
98.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	benzyl
99.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	Н	benzyl
100.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	H	Н
101.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	H
102.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	CH ₃	benzyl
103.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	Н	benzyl
104.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	Н	H
105.	4-cyclopropylnaphth-1-yl	SO ₂ NH ₂	CH ₃	H
106.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH ₃	benzyl
107.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	H	benzyl
108.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	Н	H
109.	4-cyclopropylnaphth-l-yl	F	CH ₃	H
110.	4-cyclopropylnaphth-l-yl	F	CH ₃	benzyl
111.	4-cyclopropylnaphth-l-yl	F	H	benzyl
112.	4-cyclopropylnaphth-l-yl	F	H	H
113.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	H
114.	o,o'-di-CH ₃ O-p-CN-phenyl	SO_2NH_2	CH ₃	benzyl
115.	o,o'-di-CH ₃ O-p-CN-phenyl	SO_2NH_2	H	benzyl
116.	o,o'-di-CH ₃ O-p-CN-phenyl	SO_2NH_2	Н	H
117.	o,o'-di-CH ₃ O-p-CN-phenyl	F	$ m CH_3$	H
118.	o,o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	benzyl
119.	o,o'-di-CH ₃ O-p-CN-phenyl	F	H	benzyl
120.	o,o'-di-CH ₃ O-p-CN-phenyl	F	Н	Н
121.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	CH ₃	H
122.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	CH ₃	benzyl
123.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	H	benzyl
124.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	H	Н
125.	o,o'-di-CH ₃ -p-CN-phenyl	F	CH ₃	H
126.	o,o'-di-CH ₃ -p-CN-phenyl	F	CH ₃	benzyl
127.	o,o'-di-CH ₃ -p-CN-phenyl	F	H	benzyl
128.	o,o'-di-CH ₃ -p-CN-phenyl	F	Н	H
129.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	F	H
130.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	F	benzyl

T T	Ar	V	W	Z
131.	o, o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	F	benzyl
132.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	F	H
133.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	F	H
134.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	F	benzyl
135.	4-cyclopropylnaphth-l-yl	F	F	H
136.	4-cyclopropylnaphth-l-yl	F	F	benzyl
137.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	F	H
138.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	F	benzyl
139.	o,o'-di-CH ₃ O-p-CN-phenyl	F	F	Н
140.	o,o'-di-CH ₃ O-p-CN-phenyl	F	F	benzyl
141.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	F	H
142.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	F	benzyl
143.	o,o'-di-CH ₃ -p-CN-phenyl	F	F	Н
144.	o,o'-di-CH ₃ -p-CN-phenyl	F	F	benzyl
145.	4-cyclopropyl phenyl	CN	CH ₃	H
146.	4-cyclopropyl phenyl	CN	CH ₃	benzyl
147.	4-cyclopropyl phenyl	CN	H	benzyl
148.	2,4,6-trimethyl phenyl	CN	Н	Н
149.	2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	Н
150.	2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	benzyl
151.	2,4,6-trimethyl phenyl	CH=CHCN	H	benzyl
152.	2,4,6-trimethyl phenyl	CH=CHCN	H	H
153.	2,4,6-trimethyl phenyl	CN	F	H
154.	2,4,6-trimethyl phenyl	CN	F	benzyl
155.	2,4,6-trimethyl phenyl	CH=CHCN	F	H
156.	2,4,6-trimethyl phenyl	CH=CHCN	F	benzyl
157.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	H
158.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	benzyl
159.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	H	benzyl
160.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	H	H
161.	2,4,6-trimethyl phenyl	F	CH ₃	Н
162.	2,4,6-trimethyl phenyl	F	CH ₃	benzyl
163.	2,4,6-trimethyl phenyl	F	H	benzyl
164.	2,4,6-trimethyl phenyl	F	H	H
165.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	F	H
166.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	F	benzyl
167.	4-cyclopropyl phenyl	F	F	H
168.	4-cyclopropyl phenyl	F	F	benzyl
169.	2,4,6-trimethyl phenyl	CN	CH ₃	H
170.	2,4,6-trimethyl phenyl	CN	CH ₃	benzyl
171.	2,4,6-trimethyl phenyl	CN	H	benzyl
172.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	H	H
173.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	H
174.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	benzyl
175.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	<u>H</u>	benzyl
176.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	H	H
177.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	F	H

Т	Ar	. , V	W	Z
178.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	F	benzyl
179.	o, o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	F	H
180.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	F	benzyl
181.	o, o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	CH ₃	H
182.	o, o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	CH ₃	benzyl
183.	o, o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	H	benzyl
184.	o, o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	H	H
185.	o, o'-dimethyl-p-cyclopropyl phenyl	F	CH ₃	H
186.	o, o'-dimethyl-p-cyclopropyl phenyl	F	CH ₃	benzyl
187.	o, o'-dimethyl-p-cyclopropyl phenyl	F	. H	benzyl
188.	o,o'-dimethyl-p-cyclopropyl phenyl	F	H	Н
189.	o, o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	F	H
190.	o, o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	F	benzyl
191.	2,4,6-trimethyl phenyl	F	F	H
192.	2,4,6-trimethyl phenyl	F	F	benzyl
193.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	CH ₃	CH ₃
194.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	CH ₃	cyclopropyl
195.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	H	cyclopropyl
196.	o, o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	H	CH ₃
197.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	CH ₃	CH ₃
198.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	CH ₃	cyclopropyl
199.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	H	cyclopropyl
200.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	Н	CH ₃
201.	4-cyclopropylnaphth-l-yl	CN	CH ₃	CH ₃
202.	4-cyclopropylnaphth-l-yl	CN	CH ₃	cyclopropyl
203.	4-cyclopropylnaphth-l-yl	CN	H	cyclopropyl
204.	4-cyclopropylnaphth-l-yl	CN	H	CH ₃
205.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	CH ₃
206.	4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	cyclopropyl
207.	4-cyclopropylnaphth-l-yl	CH=CHCN_	HH	cyclopropyl
208.	4-cyclopropylnaphth-l-yl	CH=CHCN	H	CH ₃
209.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	CH ₃
210.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	cyclopropyl
211.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	H	cyclopropyl
212.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	Н	CH ₃
213.	o,o'-di-CH₃O-p-CN-phenyl	CH=CHCN	CH ₃	CH ₃
214.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	cyclopropyl
215.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	H	cyclopropyl
216.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	H	CH ₃
217.	o,o'-di-CH ₃ -p-CN-phenyl	CN	CH ₃	CH ₃
218.	o,o'-di-CH ₃ -p-CN-phenyl	CN	CH ₃	cyclopropyl
219.	o,o'-di-CH ₃ -p-CN-phenyl	CN	H	cyclopropyl
220.	o,o'-di-CH ₃ -p-CN-phenyl	CN	Н	CH ₃
221.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃	CH ₃
222.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃	cyclopropyl
223.	o,o'-di-CH3-p-CN-phenyl	CH=CHCN	H	cyclopropyl
224.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	H	CH ₃

	Ar	V	W	Z
225.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	F	CH ₃
226.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CN	F	cyclopropyl
227.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	F	cyclopropyl
228.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	CH=CHCN	F	CH ₃
229.	4-cyclopropylnaphth-l-yl	CN	F	CH ₃
230.	4-cyclopropylnaphth-l-yl	CN	F	cyclopropyl
231.	4-cyclopropylnaphth-l-yl	CH=CHCN	F	CH ₃
232.	4-cyclopropylnaphth-l-yl	CH=CHCN	F	cyclopropyl
233.	o.o'-di-CH ₃ O-p-CN-phenyl	CN	F	CH ₃
234.	o,o'-di-CH ₃ O-p-CN-phenyl	CN	F	cyclopropyl
235.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	F	CH ₃
236.	o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	F	cyclopropyl
237.	o, o'-di-CH ₃ -p-CN-phenyl	CN	F	CH ₃
238.	o,o'-di-CH ₃ -p-CN-phenyl	CN	F	cyclopropyl
239.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	F	CH ₃
240.	o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	F	cyclopropyl
241.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	CH ₃
242.	o, o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	cyclopropyl
243.	o, o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	H	cyclopropyl
244.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	H	CH ₃
245.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	CH ₃	CH ₃
246.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	CH ₃	cyclopropyl
247.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	Н	cyclopropyl
248.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	Н	CH ₃
249.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH ₃	CH ₃
250.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH ₃	cyclopropyl
251.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	H	cyclopropyl
252.	4-cyclopropylnaphth-1-yl	SO_2NH_2	H	CH ₃
253.	4-cyclopropylnaphth-l-yl	F	CH ₃	CH ₃
254.	4-cyclopropylnaphth-l-yl	F	CH ₃	cyclopropyl
255.	4-cyclopropylnaphth-l-yl	F	H	cyclopropyl
256.	4-cyclopropylnaphth-l-yl	F	H	CH ₃
257.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	CH ₃
258.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	cyclopropyl
259.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	H	cyclopropyl
260.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	H	CH ₃
261.	o,o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	CH ₃
262.	o,o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	cyclopropyl
263.	o,o'-di-CH ₃ O-p-CN-phenyl	F	H	cyclopropyl
264.	o,o'-di-CH ₃ O-p-CN-phenyl	F	H	CH ₃
265.	o,o'-di-CH ₃ -p-CN-phenyl	SO_2NH_2	CH ₃	CH₃
266.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	CH ₃	cyclopropyl
267.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	H	cyclopropyl
268.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	H	CH ₃
269.	o,o'-di-CH ₃ -p-CN-phenyl	F	CH ₃	CH ₃
270.	o,o'-di-CH ₃ -p-CN-phenyl	F	CH ₃	cyclopropyl
271.	o,o'-di-CH ₃ -p-CN-phenyl	F	H	cyclopropyl

	Ar	V	W	Z
272.	o,o'-di-CH ₃ -p-CN-phenyl	F	H	CH ₃
273.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	F	CH ₃
274.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	SO ₂ NH ₂	F	cyclopropyl
275.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	F	cyclopropyl
276.	o,o'-diCH ₃ O-p-(CH=CHCN)-phenyl	F	F	CH ₃
277.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	F	CH ₃
278.	4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	F	cyclopropyl
279.	4-cyclopropylnaphth-l-yl	F	F	CH ₃
280.	4-cyclopropylnaphth-l-yl	F	F	cyclopropyl
281.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	F	CH ₃
282.	o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	F	cyclopropyl
283.	o,o'-di-CH ₃ O-p-CN-phenyl	F	F	CH ₃
284.	o,o'-di-CH ₃ O-p-CN-phenyl	F	F	cyclopropyl
285.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	F	CH ₃
286.	o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	F	cyclopropyl
287.	o,o'-di-CH ₃ -p-CN-phenyl	F	F	CH ₃
288.	o,o'-di-CH ₃ -p-CN-phenyl		F	cyclopropyl
289.	2,4,6-trimethyl phenyl	CN	CH ₃	CH ₃
290.	2,4,6-trimethyl phenyl	CN	CH ₃	cyclopropyl
291.	2,4,6-trimethyl phenyl	CN	H	cyclopropyl
292.	2,4,6-trimethyl phenyl	CN	H	CH ₃
293.	2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	CH ₃
294.	2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	cyclopropyl
295.	2,4,6-trimethyl phenyl	CH=CHCN	Н	cyclopropyl
296.	2,4,6-trimethyl phenyl	CH=CHCN	Н	CH ₃
297.	2,4,6-trimethyl phenyl	CN	F	CH ₃
298.	2,4,6-trimethyl phenyl	CN	F	cyclopropyl
299.	2,4,6-trimethyl phenyl	CH=CHCN	F	CH ₃
300.	2,4,6-trimethyl phenyl	CH=CHCN	F	cyclopropyl
301.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	CH ₃
302.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	cyclopropyl
303.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	H	cyclopropyl
304.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	Н	CH₃
305.	2,4,6-trimethyl phenyl	F	CH ₃	CH ₃
306.	2,4,6-trimethyl phenyl	F	CH ₃	cyclopropyl
307.	2,4,6-trimethyl phenyl	F	Н	cyclopropyl
308.	2,4,6-trimethyl phenyl	F	Н	CH ₃
309.	4-cyclopropyl phenyl	SO ₂ NH ₂	F	CH ₃
310.	4-cyclopropyl phenyl	SO ₂ NH ₂	F	cyclopropyl
311.	4-cyclopropyl phenyl	F	F	CH ₃
312.	4-cyclopropyl phenyl	F	F	cyclopropyl
313.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	CH ₃	CH ₃
314.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	CH ₃	cyclopropyl
315.	o, o'-dimethyl-p-cyclopropyl phenyl	CN	H	cyclopropyl
316.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	H	CH ₃
317.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	CH ₃
318.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	cyclopropyl

	Ar	V	W	Z
319.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	Н	cyclopropyl
320.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	H	CH ₃
321.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	F	CH ₃
322.	o,o'-dimethyl-p-cyclopropyl phenyl	CN	F	cyclopropyl
323.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	F	CH ₃
324.	o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	F	cyclopropyl
325.	o,o'-dimethyl-p-cyclopropyl phenyl	SO_2NH_2	CH ₃	CH ₃
326.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	CH ₃	cyclopropyl
327.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	H	cyclopropyl
328.	o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	H	CH ₃
329.	o,o'-dimethyl-p-cyclopropyl phenyl	F	CH ₃	CH ₃
330.	o,o'-dimethyl-p-cyclopropyl phenyl	F	CH ₃	cyclopropyl
331.	o,o'-dimethyl-p-cyclopropyl phenyl	F	H	cyclopropyl
332.	o,o'-dimethyl-p-cyclopropyl phenyl	F	H	CH ₃
333.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	F	CH ₃
334.	2,4,6-trimethyl phenyl	SO ₂ NH ₂	F	cyclopropyl
335.	2,4,6-trimethyl phenyl	F	F	CH ₃
336.	2,4,6-trimethyl phenyl	F	F	cyclopropyl
337.	o,o'-di-CH ₃ -p-acetyl-phenyl	CN	H	H
338.	o,o'-di-CH ₃ -p-acetyl-phenyl	CN .	CH ₃	Н
339.	o,o'-di-CH ₃ -p-acetyl-phenyl	CN	H	Cl
340.	o,o'-di-CH ₃ -p-acetyl-phenyl	CN	CH ₃	Cl

Additional contemplated and prophetic examples, which are not exhaustive but merely representative of this invention, are shown below:

Claims

What is claimed is

5 1. A compound of formula

ΙA

where the dashed line represents a double bond between either A and B or B and D, where T' is O or S; A is -N=, NZ or =CZ;

B is =CH or =N-; D is =CW or =N-, or NW, provided that at least one of A and D is -N= or NZ or NW, and further provided that when both A and D are nitrogen, then B is CH; Z is H, F, Cl, Br, CH₃, CH₂CH₃, cyclopropyl, or benzyl, in which the phenyl moiety of the benzyl group is optionally substituted with methyl or methoxy, provided that Z is not F or Cl when A is NZ;

15

20

Z is H, F, Cl, Br, CH₃, CH₂CH₃, cyclopropyl, or benzyl, the phenyl moiety of said benzyl optionally substituted with one or two groups selected independently from methyl and methoxy, provided that when A is NZ, Z is neither F nor Cl; W is H, F, Cl, Br, methyl, ethyl, cyclopropyl, allyl, CH₂CF₃, cyanomethyl, CH₂CH₂CN, CH=CHCN, or benzyl, the phenyl moiety of said benzyl optionally substituted with one or two groups selected independently from methyl and methoxy, provided that when D is NW, W is neither F or Cl; V is F, Cl, CN, SO₂CH₃, SO₂NH₂, SO₂NHCH₃, C=CCH₃, or CH=CHCN;

25

and Ar is one of (a), (b), (c), and (d) below

(a)
$$R^6$$
 R^5

(b)
$$R^6 \longrightarrow R^5$$

(c)
$$R^6 = \frac{1}{R^9} R^7$$

(d)
$$R^6 \xrightarrow{R^4} R^6$$

where the dashed lines in (a) represent optional double bonds; where R^p is Cl, Br, I, CN, methyl, ethyl, n-propyl, isopropyl, cyclopropylmethyl, C₃.C₆ cycloalkyl, CH=CHCN, acetyl, or NH-C₁-C₆ alkyl, said alkyl and cycloalkyl groups optionally substituted with methyl, methoxy, halogen, or cyano; R⁴, R⁵, and R⁶ are, independently, H, F, Cl, Br, CH₃, CH₂F, CHF₂, CF₃, isopropyl, cyclopropyl, OCH₃, OH, OCF₃, NH₂ and NHCH₃, or R⁶ and R^p on adjacent ring atoms, together with the ring atoms to which they are attached, form an additional fused five-membered ring; Q and Q' are, independently, N or CH; R⁷ is Cl, Br, I, CH₃, CF₃, OCH₃, isopropyl, cyclopropyl, *t*-butyl, or cyclobutyl; and R⁸ – R¹¹ are, independently, H or CH₃,

with the proviso that when Ar is (c), and the A,B,D ring is imidazolo, R^p and V are not both one of CH₃, CN, and CH=CHCN.

- 15 2. The compound of claim 1, wherein Ar is (a) or (c).
 - 3. The compound of claim 2, where R⁶ either is H or is in the 2-position
- 4. The compound of claim 3, wherein Ar is selected from 4-cyclopropyl phenyl; 4cyclopropylmethyl phenyl; 4-bromophenyl; 2-chloro-4-bromophenyl; 4-bromo-1-naphthyl;
 4-cyclopropyl-1-naphthyl; 2,6-dimethyl-4-cyanophenyl; 2,6-dimethoxy-4-cyanophenyl; 2,6dimethyl-4-(2-cyanoethenyl) phenyl; 2,6-dimethoxy-4-(2-cyanoethenyl) phenyl; 2-methyl-4cyclopropyl phenyl; 2,6-dimethyl-4-cyclopropyl phenyl; 2,6-di-trifluoromethyl-4-cyclopropyl
 phenyl; 2,4,6-trimethyl phenyl; and 2,6-dimethyl-4-acetyl phenyl.
- 255. The compound of claim 1, which is a compound of formula IA-1

5

10

6. The compound of claim 1, which is a compound of formula IA-2

IA-2

7. The compound of claim 1, which is a compound of formula IA-3

15 8. The compound of claim 1, which is a compound of formula IA-4

- 9. The compound of any of claims 5-8, where Ar is 4-cyclopropyl-, 4-acetyl-, 4-methyl-, 4-bromo-, or 4-cyano-2,6-di-substituted phenyl.
 - 10. The compound of claim 9, where V is CN, and where W and Z are, independently, H, methyl, halo, or benzyl.

11. The compound of claim 3 which is a compound selected from the two structures below

$$\mathbb{R}^6$$
 \mathbb{R}^7
 \mathbb{R}^7

12. The compound of claim 11, where V is CN or CH=CHCN, R⁶ is 2-methyl, 2-methoxy, or

- 13. The compound of claim 12, where R^p is CN, cyclopropyl, methyl, Br, Cl, CH=CHCN, or
- 15 14. The compound of claim 13 which is selected from the group consisting of

2-chloro and R⁷ is H, 6-methyl, or 6-methoxy.

- 1) a compound of structure IA-1a, where V is CN, W is H, methyl, ethyl, or benzyl, and Z is H, chloro, bromo, methyl, or ethyl; and
- 2) a compound of structure IA-2a, where V is CN, W is H, chloro, bromo, methyl, or ethyl and Z is H, methyl, ethyl, or benzyl.

20

5

10

acetyl.

15. A compound of formula IB

25

where the dashed line represents a double bond between either A and B or B and D, and where A is -N=, NZ, or CZ;

B is CH or =N-

D is CW or =N-, or NW, provided that at least one of A and D is -N= or NZ or NW, and further provided that when both A and D are nitrogen, then B is CH;

- Z is H, F, Cl, Br, CH₃, CH₂CH₃, cyclopropyl, or benzyl, the phenyl moiety of said benzyl optionally substituted with methyl or methoxy, provided that when A is NZ, Z is neither F nor Cl; W is H, F, Cl, Br, methyl, ethyl, cyclopropyl, allyl, CH₂CF₃, cyanomethyl, CH₂CH₂CN, CH=CHCN, or benzyl, the phenyl moiety of said benzyl optionally substituted with methyl or methoxy, provided that when D is NW, W is neither F or Cl;
- V is F, Cl, CN, SO₂CH₃, SO₂NH₂, SO₂NHCH₃, C≡CCH₃, or CH=CHCN, and Ar is one of (a), (b), and (d) below:

$$R^{6} \xrightarrow{R^{5}} R^{2}$$

$$R^{5} \xrightarrow{R^{5}} R^{5}$$

$$R^{6} \xrightarrow{R^{5}} R^{5}$$

$$R^{8} \xrightarrow{R^{10}} R^{11}$$

$$R^{11} (d)$$

15

where R^p is Cl, Br, I, CN, methyl, ethyl, n-propyl, isopropyl, cyclopropylmethyl, C₃-C₆ cycloalkyl, CH=CHCN, acetyl, or NH-C₁-C₆ alkyl, said alkyl and cycloalkyl groups optionally substituted with methyl, methoxy, halogen, or cyano; R⁴, R⁵, and R⁶ are, independently, H, F, Cl, Br, CH₃, CH₂F, CHF₂, CF₃, isopropyl, cyclopropyl, OCH₃, OH, OCF₃, NH₂ and NHCH₃; Q and Q' are, independently, N or CH; R⁷ is Cl, Br, I, CH₃, CF₃, OCH₃, isopropyl, cyclopropyl, *t*-butyl, or cyclobutyl; and R⁸ – R¹¹ are, independently, H or CH₃.

16. The compound of claim 14 which is a compound of formula IB-1

25

20

30

IB-1

where the dashed line represents a double bond between either A and B or B and D; where A is -N=, NZ or CZ;

- 5 B is CH or =N-
 - D is CW or =N-, or NW;
 - Z is H, F, Cl, Br, CH₃, CH₂CH₃, cyclopropyl, or benzyl, the phenyl moiety of said benzyl optionally substituted with methyl or methoxy, provided that when A is NZ, Z is neither F nor Cl;
- W is H, F, Cl, Br, methyl, ethyl, cyclopropyl, allyl, CH₂CF₃, CH₂CN, CH₂CH₂CN, CH=CHCN, or benzyl, the phenyl moiety of said benzyl optionally substituted with methyl or methoxy, provided that when D is NW, W is neither F nor Cl;
 V is F, Cl, CN, SO₂CH₃, SO₂NH₂, SO₂NHCH₃, C=CCH₃, or CH=CHCN;
 provided that at least one of A and D is -N= or NZ or NW, and further provided that when
 both A and D are nitrogen, then B is CH and V is other than CN or CH=CHCN;
 - wherein R^p is Cl, Br, I, CN, CH=CHCN, methyl, ethyl, n-propyl, isopropyl, cyclopropylmethyl, C₃-C₆ cycloalkyl, acetyl, and NH-C₃-C₆ alkyl; R⁶ is H, F, Cl, Br, CH₃, CH₂F, CHF₂, CF₃, isopropyl, cyclopropyl, OCH₃, OH, OCF₃, NH₂ and NHCH₃; R⁷ is Cl, Br,
- 20 I, CH₃, CF₃, OCH₃, isopropyl, cyclopropyl, *t*-butyl, or cyclobutyl.
 - 17. The compound of claim 16, where V is CN or CH=CHCN, R⁶ is 2-methyl, 2-methoxy, or 2-chloro and R⁷ is H, 6-methyl, or 6-methoxy.
- 25 18. The compound of claim 17, where R^p is CN, cyclopropyl, methyl, Br, Cl, CH=CHCN, or acetyl.
 - 19. The compound of claim1, where one of A and D is =N- and the other is NZ or NW.
- 30 20. The compound of claim 5 which is a compound of formula IA-1 selected from compounds in Table 1:

T	٨		1
	А	_	1

Ar	V	W	Z
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	H	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	benzyl	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	benzyl	Н
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	3-Me-benzyl	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	4-Me-benzyl	H
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	3-MeO-benzyl	H
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	4-MeO-benzyl	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	H	H
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	Н	Br
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	cyclopropyl	CH ₂ CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	CH ₂ CF ₃	CH ₂ CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	Н	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	benzyl	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	benzyl	Н
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	3-Me-benzyl	cyclopropyl
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	3-MeO-benzyl	benzyl
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	Н	Н
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	C≡CCH ₃	CH ₂ CH ₃	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	Cl	CH ₂ CH=CH ₂	H
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ CH ₃	CH ₂ CH=CH ₂	H
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	C1	CH ₂ CH ₃	CH₂CH₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	Cl	H	H
4-cyclopropylnaphth-l-yl	CN	H	CH ₃
4-cyclopropylnaphth-l-yl	CN	benzyl	CH_3
4-cyclopropylnaphth-l-yl	CN	benzyl	H
4-cyclopropylnaphth-l-yl	CN	H	H
4-cyclopropylnaphth-l-yl	CH=CHCN	H	CH ₃
4-cyclopropylnaphth-l-yl	CH=CHCN	benzyl	CH ₃
4-cyclopropylnaphth-l-yl	CH=CHCN	benzyl	<u>H</u>
4-cyclopropylnaphth-l-yl	CH=CHCN	H	H
4-cyclopropylnaphth-l-yl	SO ₂ NHCH ₃	CH ₂ CN	F
4-cyclopropylnaphth-l-yl	SO ₂ NHCH ₃	cyclopropyl	C1
o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₂ CH ₂ CN	Br
o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₂ CN	benzyl
o,o'-di-CH ₃ O-p-CN-phenyl	C≡CCH ₃	3-MeO-benzyl	F
o,o'-di-CH ₃ O-p-CN-phenyl	F	3-Me-benzyl	C1
o,o'-di-CH ₃ O-p-CN-phenyl	CN	Н	CH ₃
o,o'-di-CH ₃ O-p-CN-phenyl	CN	benzyl	CH ₃
o,o'-di-CH ₃ O-p-CN-phenyl	CN	benzyl	H

PCT/US2006/017677

Ar	V	W	Z
o,o'-di-CH ₃ O-p-CN-phenyl	CN	H	H
o, o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	H	CH ₃
o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	benzyl	CH ₃
o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	benzyl	H
o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	H	H
o,o'-di-CH ₃ -p-CN-phenyl	CN	H	CH ₃
o,o'-di-CH ₃ -p-CN-phenyl	CN	benzyl	CH ₃
o,o'-di-CH ₃ -p-CN-phenyl	CN	3,5-di MeO-	CH ₃
		benzyl	
o,o'-di-CH ₃ -p-CN-phenyl	CN	benzyl	H
o,o'-di-CH ₃ -p-CN-phenyl	CN	Н	H
o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	Н	CH ₃
o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	benzyl	CH ₃
o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	benzyl	H
o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	H	H
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	H	F
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	benzyl	F
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	benzyl	F
o, o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	H	F
4-cyclopropylnaphth-l-yl	CN	H	F
4-cyclopropylnaphth-l-yl	CN	benzyl	F
4-cyclopropylnaphth-l-yl	CH=CHCN	H	F
4-cyclopropylnaphth-l-yl	CH=CHCN	benzyl	F
o,o'-di-CH ₃ O-p-CN-phenyl	CN	H	F
o,o'-di-CH ₃ O-p-CN-phenyl	CN	benzyl	F
o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	H	F
o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	benzyl	F
o,o'-di-CH ₃ -p-CN-phenyl	CN	H	F
o,o'-di-CH ₃ -p-CN-phenyl	CN	benzyl	F
o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	H	F
o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	benzyl	F
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	H	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	benzyl	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	benzyl	H
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	H	H
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	H	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	benzyl	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	benzyl	H
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	H	H
4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	H	CH ₃
4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	benzyl	CH ₃
4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	benzyl	H
4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	H	H
4-cyclopropylnaphth-l-yl	F	H	CH ₃
4-cyclopropylnaphth-l-yl	F	benzyl	CH ₃
4-cyclopropylnaphth-l-yl	F	benzyl	Н
4-cyclopropylnaphth-l-yl	F	H	н

Ar	V	W	Z
o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	Н	CH ₃
o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	benzyl	CH ₃
o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	benzyl	H
o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	H	H
o,o'-di-CH ₃ O-p-CN-phenyl	F	Н	CH ₃
o,o'-di-CH ₃ O-p-CN-phenyl	F	benzyl	CH ₃
o,o'-di-CH ₃ O-p-CN-phenyl	F	benzyl	H
o,o'-di-CH ₃ O-p-CN-phenyl	F	Н	Н
o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	Н	CH ₃
o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	benzyl	CH ₃
o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	3-Me-benzyl	CH ₃
o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NHCH ₃	benzyl	H
o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	benzyl	H
o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	H	H
o,o'-di-CH ₃ -p-CN-phenyl	F	H	CH ₃
o,o'-di-CH ₃ -p-'CN-phenyl	F	benzyl	CH ₃ '
o,o'-di-CH ₃ -p-CN-phenyl	F	benzyl	H
o,o'-di-CH ₃ -p-CN-phenyl	F	H	Н
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	Н	F
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	benzyl	F
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	benzyl	F
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	H	F
4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	H	F
4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	benzyl	F
4-cyclopropylnaphth-l-yl	F	H	F
4-cyclopropylnaphth-l-yl	F	benzyl	F
o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	H	F
o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	benzyl	F
o,o'-di-CH ₃ O-p-CN-phenyl	F	H	F
o,o'-di-CH ₃ O-p-CN-phenyl	F	benzyl	F
o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	H	F
o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	benzyl	F
o,o'-di-CH ₃ -p-CN-phenyl	F	H	F
o,o'-di-CH ₃ -p-CN-phenyl	F	benzyl	F
2,4,6-trimethyl phenyl	CN	H	CH ₃
2,4,6-trimethyl phenyl	CN	benzyl	CH ₃
2,4,6-trimethyl phenyl	CN	benzyl	H
2,4,6-trimethyl phenyl	CN	H	Н
2,4,6-trimethyl phenyl	CH=CHCN	H	CH ₃
2,4,6-trimethyl phenyl	CH=CHCN	benzyl	CH ₃
2,4,6-trimethyl phenyl	CH=CHCN	benzyl	H
2,4,6-trimethyl phenyl	CH=CHCN	H	H
2,4,6-trimethyl phenyl	CN	H	F
2,4,6-trimethyl phenyl	CN	benzyl	F
2,4,6-trimethyl phenyl	CH=CHCN	H	F
2,4,6-trimethyl phenyl	CH=CHCN	benzyl	F
2,4,6-trimethyl phenyl	SO ₂ NH ₂	H	CH ₃

Ar	V	W	Z
2,4,6-trimethyl phenyl	SO ₂ NH ₂	benzy1	CH ₃
2,4,6-trimethyl phenyl	SO ₂ NH ₂	benzyl	H
2,4,6-trimethyl phenyl	SO ₂ NH ₂	H	H
2,4,6-trimethyl phenyl	F	H	CH ₃
2,4,6-trimethyl phenyl	F	benzyl	CH ₃
2,4,6-trimethyl phenyl	F	benzyl	H
4-cyclopropyl phenyl	F	H	H
4-cyclopropyl phenyl	SO ₂ NH ₂	H	F
4-cyclopropyl phenyl	SO ₂ NH ₂	benzyl	F
4-cyclopropyl phenyl	F	H	F ·
4-cyclopropyl phenyl	F	benzyl	F
o,o'-dimethyl-p-cyclopropyl phenyl	CN	H	CH ₃
o,o'-dimethyl-p-cyclopropyl phenyl	CN	benzyl	CH ₃
o,o'-dimethyl-p-cyclopropyl phenyl	CN	benzyl	H
o,o'-dimethyl-p-cyclopropyl phenyl	CN	H	H
o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	H	CH ₃
o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	benzyl	CH ₃
o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	benzyl	H
o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	Н	H
o,o'-dimethyl-p-cyclopropyl phenyl	CN	H	F
o,o'-dimethyl-p-cyclopropyl phenyl	CN	benzyl	F
o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	Н	F
o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	benzyl	F
o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	Н	CH ₃
o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	benzyl	CH ₃
o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	benzyl	H
o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	Н	H
o,o'-dimethyl-p-cyclopropyl phenyl	F	H	CH ₃
o,o'-dimethyl-p-cyclopropyl phenyl	F	benzyl	CH₃
o,o'-dimethyl-p-cyclopropyl phenyl	F	benzyl	H
o,o'-dimethyl-p-cyclopropyl phenyl	F	H	H
o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	H	F
o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	benzyl	F
o,o'-dimethyl-p-cyclopropyl phenyl	F	H	F
o,o'-dimethyl-p-cyclopropyl phenyl	F	benzyl	F
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	CH ₃	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	cyclopropyl	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	cyclopropyl	<u>H</u>
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	CH ₃	H
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	CH ₃	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	cyclopropyl	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	cyclopropyl	H
o, o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	CH ₃	H
4-cyclopropylnaphth-l-yl	CN	CH ₃	CH ₃
4-cyclopropylnaphth-l-yl	CN	cyclopropyl	CH ₃
4-cyclopropylnaphth-l-yl	CN	cyclopropyl	Н
4-cyclopropylnaphth-l-yl	CN	CH ₃	H

Ar	V	W	Z
4-cyclopropylnaphth-1-yl	CH=CHCN	CH ₃	CH ₃
4-cyclopropylnaphth-l-yl	CH=CHCN	cyclopropyl	CH ₃
4-cyclopropylnaphth-l-yl	CH=CHCN	cyclopropyl	H
4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	H
o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	CH ₃
o,o'-di-CH ₃ O-p-CN-phenyl	CN	cyclopropyl	CH ₃
o,o'-di-CH ₃ O-p-CN-phenyl	CN	cyclopropyl	H
o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	H
o,o'-di-CH₃O-p-CN-phenyl	CH=CHCN	CH ₃	CH ₃
o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	cyclopropyl	CH ₃
o,o'-di-CH₃O-p-CN-phenyl	CH=CHCN	cyclopropyl	H
o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	Н
o,o'-di-CH ₃ -p-CN-phenyl	CN	CH ₃	CH ₃
o,o'-di-CH ₃ -p-CN-phenyl	CN	cyclopropyl	CH ₃
o,o'-di-CH ₃ -p-CN-phenyl	CN	cyclopropyl	H
o,o'-di-CH ₃ -p-CN-phenyl	CN	CH ₃	Н .
o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃	CH ₃
o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	cyclopropyl	CH ₃
o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	cyclopropyl	H
o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃ .	H
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	CH ₃	F
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CN	cyclopropyl	F
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	cyclopropyl	F
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	CH=CHCN	CH ₃	F
4-cyclopropylnaphth-l-yl	CN	CH ₃	F
4-cyclopropylnaphth-l-yl	CN	cyclopropyl	F
4-cyclopropylnaphth-l-yl	CH=CHCN	CH ₃	F
4-cyclopropylnaphth-l-yl	CH=CHCN	cyclopropyl	F
o,o'-di-CH ₃ O-p-CN-phenyl	CN	CH ₃	F
o,o'-di-CH ₃ O-p-CN-phenyl	CN	cyclopropyl	F
o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	CH ₃	F
o,o'-di-CH ₃ O-p-CN-phenyl	CH=CHCN	cyclopropyl	F
o,o'-di-CH ₃ -p-CN-phenyl	CN	CH ₃	F
o,o'-di-CH ₃ -p-CN-phenyl	CN	cyclopropyl	F
o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	CH ₃	F
o,o'-di-CH ₃ -p-CN-phenyl	CH=CHCN	cyclopropyl	F
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	CH ₃	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	cyclopropyl	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	cyclopropyl	H
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	CH ₃	Н
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NHCH ₃	CH ₃	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	cyclopropyl	CH ₃
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	cyclopropyl	H
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	CH ₃	H
4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH ₃	CH ₃
4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	cyclopropyl	CH ₃
4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	cyclopropyl	H

Ar	V	W	Z
4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH₃	H
4-cyclopropylnaphth-l-yl	F	CH ₃	CH ₃
4-cyclopropylnaphth-l-yl	F	cyclopropyl	CH ₃
4-cyclopropylnaphth-l-yl	F	cyclopropyl	H
4-cyclopropylnaphth-l-yl	F	CH ₃	H
o, o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	CH ₃
o, o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NHCH ₃	CH ₃	CH ₃
o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	CH ₃
o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	H
o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	H
o, o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	CH ₃
o,o'-di-CH ₃ O-p-CN-phenyl	F	cyclopropyl	CH ₃
o,o'-di-CH ₃ O-p-CN-phenyl	F	cyclopropyl	H
o,o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	H
o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	CH ₃	CH ₃
o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	CH ₃
o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	H
o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	CH ₃	H
o,o'-di-CH ₃ -p-CN-phenyl	F	CH ₃	CH ₃
o,o'-di-CH ₃ -p-CN-phenyl	F	cyclopropyl	CH ₃
o,o'-di-CH ₃ -p-CN-phenyl	F	cyclopropyl	H
o,o'-di-CH ₃ -p-CN-phenyl	F	CH ₃	H
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	CH ₃	F
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	SO ₂ NH ₂	cyclopropyl	F
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	cyclopropyl	F
o,o'-diCH ₃ O-p-(CH=CHCN)phenyl	F	CH ₃	F
4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	CH ₃	F
4-cyclopropylnaphth-l-yl	SO ₂ NH ₂	cyclopropyl	F
4-cyclopropylnaphth-l-yl	F	CH ₃	F
4-cyclopropylnaphth-l-yl	F	cyclopropyl	F
o,o'-di-CH₃O-p-CN-phenyl	SO ₂ NH ₂	CH ₃	F
o,o'-di-CH ₃ O-p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	F
o, o'-di-CH ₃ O-p-CN-phenyl	F	CH ₃	F
o, o'-di-CH ₃ O-p-CN-phenyl	F	cyclopropyl	F
o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	CH ₃	F
o,o'-di-CH ₃ -p-CN-phenyl	SO ₂ NH ₂	cyclopropyl	F
o, o'-di-CH ₃ -p-CN-phenyl	F	CH ₃	F
o,o'-di-CH ₃ -p-CN-phenyl	F	cyclopropyl	F
4-cyclopropyl phenyl	CN	CH ₃	CH ₃
2,4,6-trimethyl phenyl	CN	cyclopropyl	CH ₃
2,4,6-trimethyl phenyl	CN	cyclopropyl	<u>H</u>
2,4,6-trimethyl phenyl	CN	CH ₃	<u>H</u>
2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	CH ₃
2,4,6-trimethyl phenyl	CH=CHCN	cyclopropyl	CH ₃
2,4,6-trimethyl phenyl	CH=CHCN	cyclopropyl	H
2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	H
2,4,6-trimethyl phenyl	CN	CH ₃	F

Ar	V	W	Z
2,4,6-trimethyl phenyl	CN	cyclopropyl	F
2,4,6-trimethyl phenyl	CH=CHCN	CH ₃	F
2,4,6-trimethyl phenyl	CH=CHCN	cyclopropyl	F
2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	CH ₃
2,4,6-trimethyl phenyl	SO ₂ NH ₂	cyclopropyl	CH ₃
2,4,6-trimethyl phenyl	SO ₂ NH ₂	cyclopropyl	H
2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	H
2,4,6-trimethyl phenyl	F	CH ₃	CH ₃
2,4,6-trimethyl phenyl	F	cyclopropyl	CH ₃
2,4,6-trimethyl phenyl	F	cyclopropyl	H
4-cyclopropyl phenyl	F	CH ₃	H
4-cyclopropyl phenyl	SO ₂ NH ₂	CH ₃	F
4-cyclopropyl phenyl	SO ₂ NH ₂	cyclopropyl	F
4-cyclopropyl phenyl	F	CH ₃	F
4-cyclopropyl phenyl	F	cyclopropyl	F
2,4,6-trimethyl phenyl	CN	CH ₃	CH ₃
o,o'-dimethyl-p-cyclopropyl phenyl	CN	cyclopropyl	CH ₃
o,o'-dimethyl-p-cyclopropyl phenyl	CN	cyclopropyl	Н
o,o'-dimethyl-p-cyclopropyl phenyl	CN	CH ₃	H
o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	CH ₃
o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	cyclopropyl	CH ₃
o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	cyclopropyl	H
o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	H
o,o'-dimethyl-p-cyclopropyl phenyl	CN	CH₃	F
o,o'-dimethyl-p-cyclopropyl phenyl	CN	cyclopropyl	F
o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	CH ₃	F
o,o'-dimethyl-p-cyclopropyl phenyl	CH=CHCN	cyclopropyl	F
o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	CH ₃	CH ₃
o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	cyclopropyl	CH ₃
o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	cyclopropyl	H
o,o'-dimethyl-p-cyclopropyl phenyl	SO ₂ NH ₂	CH ₃	H
o,o'-dimethyl-p-cyclopropyl phenyl	F	CH ₃	CH ₃
o,o'-dimethyl-p-cyclopropyl phenyl	F	cyclopropyl	CH ₃
o,o'-dimethyl-p-cyclopropyl phenyl	F	cyclopropyl	H
2,4,6-trimethyl phenyl	F	CH ₃	Н
2,4,6-trimethyl phenyl	SO ₂ NH ₂	CH ₃	F
2,4,6-trimethyl phenyl	SO ₂ NH ₂	cyclopropyl	F
2,4,6-trimethyl phenyl	F	CH ₃	F
2,4,6-trimethyl phenyl	F	cyclopropyl	F
o,o'-di-CH ₃ -p-acetyl-phenyl	CN	CH ₃	H
o,o'-di-CH ₃ -p-acetyl-phenyl	CN	H	H
o,o'-di-CH ₃ -p-acetyl-phenyl	CN	CH ₃	Cl
o,o'-di-CH ₃ -p-acetyl-phenyl	CN	<u> </u> H	Cl